be, the sigmoid flexure. The child having been chloro-formed, I made an incision an inch and a half in length, extending in an oblique direction upwards and outwards from the middle line to a point about half an inch beyond the anterior superior spinous process of the ilium, dividing the skin, and cellular tissue or superficial fasciæ till I reached the muscles, which with the transversalis fascia. and peritoneum were successively divided on the director. The distended bowel lay immediately beneath ready to hand. The gut was transfixed by two curved parallel needles, half an inch apart, and each threaded with double ligatures of carbolized catgut, after which it was opened and the edges attached to those of the wound in the integument. A large quantity of meconium of a greyishvellow colour was immediately voided. The substance of the coats of the bowel was found to be very easily lacerated. as if softened, from the effects perhaps of previous inflammation from the distension. The operation, although tedious, was one which presented no very great difficulties in its performance. Some bleeding from the lower extremity of the wound was arrested without much difficulty. At no time after the operation had the child a single bad symptom but nursed and slept as if nothing had happened. The skin rapidly lost the dirty sallow hue it previously presented, and at the end of a fortnight the little fellow became quite plump and healthy-looking.

Repeated explorations of the bowel, first with elastic bougies, and subsequently with a whalebone probe, introduced through the opening in the groin, conjoined with the finger in the canal cul de sac, lead to the conclusion that the rectum is absent for a portion of its length, as the probe thus used cannot be made to pass downward for more than about three quarters of an inch, and cannot be felt by the finger. The most remarkable fact in the case, perhaps the only one excepting of course the success of an operation so frequently fatal, is that this child should have lived for a fortnight (as it did) previous to the operation, in a condition of compara-