

SCAPANUS TOWNSENDI, *Bach.*

Oregon Mole.

*Scalops Canadensis*, Rich.*Scalops latimanus*, Bach.*Scalops aeneus*, Cassin.

Southern portions of British Columbia.

SCAPANUS BREWERI, *Bach.*

Hairy-tailed Mole.

New Brunswick.

CONDYLURA CRISTATA, *Linn.*

Star-nosed Mole.

*Condylura longicauda*, (Ill.) Rich.*Condylura macroura*, Rich.From the Atlantic Coast north westward to Hudson's Bay  
and westward to the Rocky Mountains.UROTRICHUS GIBBSI, *Baird.*

Chilukweynuk River in southern British Columbia.

NEOSOREX PALUSTRIS, *Rich.*

Marsh Shrew.

Nova Scotia to the Rocky Mountains around the borders of  
lakes.SOREX TROWBRIDGII, *Baird.*

Southern British Columbia.

SOREX FORSTERI, *Rich.*

Forster's Shrew.

From Nova Scotia to the Rocky Mountains, and as far north  
on the Mackenzie River as the Arctic Circle.SOREX THOMPSONI, *Baird.*

Thompson's Shrew.

Halifax, Nova Scotia.

SOREX VAGRANS, (Cooper) *Baird.*

Southern British Columbia.