

black dots; palpi orange yellow tipped with black, antennæ black. Thorax white, yellowish behind, with a broad central brown band. Beneath, thorax yellow, legs yellow, fore coxæ with a black or brown dot, fore and median femora and tibiæ dark brown exteriorly, fore and median tarsi brown. Abdomen ochre-yellow with a dorsal row of brown dots which are sometimes almost obsolete. Primaries marked exactly as in *Suffusa*, (fig. 11). Secondaries ochre-yellow, immaculate, or with a brown dot near anal angle. Beneath, all the wings are deep ochre-yellow, the brown markings of primaries being reproduced more or less distinctly except towards apex and outer margin. Expands, 53-58 mm. = 2 $\frac{1}{5}$ -2 $\frac{1}{4}$ inch.

This species is well known and distinct. It varies somewhat in the maculation of primaries, but not so far as ever to render its determination doubtful.

CALLIMORPHA INTERRUPTO-MARGINATA, De Beauv.

Bombix Interrupto-marginata, De Beauv., Ins. Afriq. et Amer.,
p. 265, pl. 24, figs. 5, 6.

Callimorpha Anchora, Harris, MS.

Hypercompa Comma, Walk. Cat. Lep. B. M. III., p. 652.

Head whitish, front and vertex ochre-yellow, palpi yellow tipped with black. Collar and thorax white with a broad median brown band. Beneath, legs as in *Clymene* except that the spot on fore coxæ is wanting. Abdomen ochre-yellow with a dorsal brown stripe.

Primaries creamy white, suffused with yellow at the apex, with a dark brown stripe along the costa to near the apex, a band of the same colour along the inner margin to inner angle, near which it branches, sending forward half way to the costa a broad, slightly curved band, which marks, when the wings are closed, have a slight resemblance to an anchor.

Outer margin with an incomplete brown border, widest at the middle.

Secondaries ochre-yellow with a brown spot near anal angle.

Beneath ochre-yellow with the brown markings partially reproduced.

Expands, 47-49 mm.

Hab., Canada to Florida.

This very distinct species seems to be the most constant of any and it is a shame that it should be burdened with such an abominable name. The markings are very close to fig. 5, except that there is no spur on the costal border, and that there is the brown spot on secondaries.