CORNED VENISON.—When you have more fresh meat of this kind than you think will keep good, rub it with salt, and hang it in the root-house or dairy.

Venison-Ham.—Make a mixture of sugar, and a very little saltpetre; rub the haunch well with this every day, for three weeks; hang it to smoke for three more. It is very good grated, or if dried, cut in thin shavings, as a relish with bread and butter for tea or breakfast, with salad. Jerked venison is the flesh cut in strips and dried in the open air.

ERYSIPELAS.—A correspondent of the Providence Journal says, that in ninetynine cases out of every hundred, cranberries applied as a poultice will effectually cure the erysipelas. There is not an instance known where it has failed to effect a cure, when faithfully applied before the sufferer was in a dying state. Two applications generally do the work.

MONTHLY METEOROLOGICAL REPORT FOR AUGUST, 1858.

From observations taken at St. Martin, Ile Jesus, C. E., Latatude 45 degrees 32 minutes, Longitude, 73 degrees, 36 minutes West, Height over the level of the sea 118 feet.

BY CHS. SMALLWOOD, M. D. L. L. D.

BAROMETER.	Greatest intensity of the suns
Mean reading of the barometer F inches	rays
corrected and reduced to 32° 29.771	tion
Highest reading of the barometer 30° 002 Lowest reading of the barometer 29° 342	Amount of evaporation in inches 3 69 Rain fell on 13 days amounting
Monthly range 0 660	to 8.656 inches it was raining
THERMOMETER.	49 hours 51 minutes, accompanied by Thunder on 4 days.
Mean reading of the standard	Most prevalent wind S. E
thermometer	Least prevalent wind N
Hightest reading of the maxi-	Most windy day the 5th, mean
mum do 97° 4	miles per hour
Lowest reading of the minimum	Least do do the 23 day do 0 00
do	Ozone was present in moderate
Monthly Range	quantity
Mean of humidity 0° 756	Aurora borealis visible on 1 night