

## ONOCLEA

sensibilis (*Sensitive Fern*).

In wet shady places. The sterile fronds extremely variable; general and very common.

## OSMUNDA (The Flowering Ferns.)

regalis (*Royal Fern*).

In moist woods and swamps; general and very common.

The American plant differs slightly from the European, and is known as var. *spectabilis*.

Claytoniana (*Interrupted Flowering-fern*).

As the last; equally general and abundant.

cinnamomea (*Cinnamon-fern*).

As the last; general and common.

## BOTRYCHIUM (The Moonworts.)

lunarioides (*Tall Moonwort*).

In dry woods and clearings. A very variable plant; generally discontinued from Quebec westward, but not plentiful.

virginicum (*Rattlesnake-fern*).

In rich woods. A variable plant; generally distributed and common.

Lunaria (*Common Moonwort*).

In open woods. A northern plant. General from Hastings (Mr. John Bell) eastwards, but apparently rare; being inconspicuous it may have escaped observation. This plant—the *B. Lunaria* (Linn.) of Swartz and all modern authors—is common in many parts of Europe. It is not in Gray's Manual.

simplex (*Dwarf Moonwort*).

In woods. As the last, of which it may possibly be only a variety.

## OPHIOGLOSSUM

vulgatum (*Adder's-tongue*).

In bogs and wet woods. General and probably not uncommon. A variable plant.

## LYCOPODIUM (The Club Mosses.)

lucidulum (*Shining Club-moss*).

In cold swampy woods. General, and usually common.

inundatum (*Marsh Club-moss*).

River Mistassini and northward, Michaux, MSS., per Abbé Brunet; Hastings, Mr. Macoun; probably general.

annotinum (*Interrupted Club-moss*).

In woods. A somewhat variable plant; general, and common.