

*Pinipestis umbripennis*, n. sp.—Expands 24 mm. Palpi black, a little grayish at tips; head and thorax black; abdomen blackish, a little lighter on each segment posteriorly. Fore wings mouse-black, slightly grayish across basal portion within scale ridge; also a triangular, dark grayish space along costa on middle field, enclosing double black discal spots, and a faint grayish marginal shade. Basal line not apparent; scale ridge strong, jet black; outer line narrow, dark gray, bent inward at end of cell, then with three even teeth outward, then angled inward before reaching inner margin. Hind wings deep black, with a blackish-brown reflection. Beneath almost uniform dull somewhat glistening black.

Colorado. One ♂, from Mr. Bruce.

*Salebria delectella*, n. sp.—Expands 33 mm. Palpi thin, lightly scaled, gray; thorax blackish-gray, with a subdorsal tuft of orange-yellow scales on each side; abdomen blackish-gray, and gray-ringed on each segment. Fore wings, general colour blackish-gray; basal space, except along costa, dull reddish; middle field, first half of ground colour, the outer half much lighter gray; outer field, except towards costa, dull reddish; discal spot a large white lunule, concave side outward; basal line light gray, well out, wavy, dentate; outer line gray, somewhat indeterminate on borders, with a more prominent outward dentation at middle; marginal line black. Hind wings light translucent fuscous, with black marginal line, except along inner margin. Beneath, fore wings dark fuscous, the outer line faintly showing; hind wings much as above.

Colorado. Two ♀♀, from Mr. Bruce.

In general appearance somewhat resembling *Salebria tarmitalis*, which is now put by Mr. Ragonot under the genus *Myrcala*. This may also belong there, as the cell of the hind wings is very short. But as I have no male, the position of the insect is in doubt. The thoracic tufts are somewhat peculiar.

*Salebria georgiella*, n. sp.—Expands 17 mm. Palpi fuscous gray, blackish in front, strong, heavily scaled, reaching above head; maxillary palpi strongly pencil tufted, bright yellow; antennæ brownish fuscous, the scale tuft in bend very heavy; head fuscous gray; thorax gray or bluish-gray; abdomen ringed with ochre-fuscous and yellow ochre, somewhat tufted at end. Fore wings gray, washed with fuscous along costa, becoming clear gray posteriorly; a dull brick-red central dash at base, and a large dull brick-red spot within basal line posterior to centre and reaching