

2026. *Homalota maritima*, Mann.—Common. Frequents stones near low-water mark on beach.
2027. *Homalota picipennis*, Mann.—Numerous in June, under excrement.
2033. *Homalota geniculata* Makl.—Common on beach in loose dry sand near high-water mark. Under driftwood. Shaken from rotten sea-weed, June 3. Emerged and took flight on sand being pressed, April and May.
2036. *Homalota comparabilis*, Makl.—One taken in April, under driftwood on sandy beach.
2039. *Homalota fucicola*, Makl.—June 3; in rotten sea-weed.
- 2040½. *Homalota atricornis*, Fol.—Not common. Under excrement in August.
- Homalota fungi*, Grav.—Common under excrement during summer; rotten grass in June; occasionally in turf in winter.
- Homalota ingrata*, Fvl.—Numerous under carrion in May. Jet black. Legs dusky yellow.
- Aleochara californica*, Fvl.
2056. *Aleochara castaneipennis*, Mann., var. with rufous elytra.—Common under carrion, all summer.
2057. *Aleochara sulcicollis*, Mann.—Numerous under rotten sea-weed.
2058. *Aleochara cognata*, Makl.
- Aleochara littoralis*, Makl.—Under driftwood on beach.
- 2064½. *Oxypoda crassicornis*, Fvl.—January 4. Common in flood refuse.
9292. *Autalia elegans*, Casey.—Numerous in August.
- Bryobiotus Keeni*, Fvl., n. sp.—Occasional in June, under stones on sandy beach, between tide marks. Larvæ in same place.
- Phytosus Fletcheri*, Fvl.—Rare; taken in May under driftwood.
2071. *Bolitochara notata*, Makl.—(= *Californica*, Casey).—Occasional; under bark. One in June, 1893, in rotten grass. Three on carrion in March.
2075. *Leptusa*.—Occasionally under bark. March to October.
- Heterothops asperatus*, Fvl., n. sp.—Several taken in April, under driftwood on sandy beach.
9479. *Quedius erythrogaster*, Mann.—Rare; in manure heaps in May and June.
2101. *Quedius sublimbatus*, Makl.—Occasional; in moss in winter.
- Quedius marginalis*, Makl.—Occurs occasionally in summer in rotten grass.