

(6.) *E. MINUTUM*, Lindl.

From several localities in British Columbia and common on Vancouver Island. The form named *adscendens* by Suksdorf, was collected on Mt. Benson, Van. Island, by Prof. Macoun in 1893.

Var. *FOLIOSUM*, Torr & Gray.

Sproat, Columbia River, B.C., (*John Macoun*) and Yale Mt., B.C., (*Fletcher*.) Common on Vancouver Island.

(7.) *E. STRICTUM*, Muhl.

*E. molle*, Torr. of Macoun Cat. of Canadian Plants, p. 171 in part and p. 530.

Specimens in our herbarium are from East Pt., P.E.I., and Belleville, Ont. (*John Macoun*); Cartwright, Ont. (*W. Scott*.)

(8.) *E. LINEARE*, Muhl.

Common from Prince Edward Island west to Beaver Creek, Selkirk Mts., B.C.

(9.) *E. PALUSTRE*, L.

Common from the Atlantic to the Rocky Mts. No specimens in our herbarium from British Columbia, but found north of that province by Dr. G. M. Dawson on the Lewis River in Lat. 62°.

(10.) *E. PALUSTRE* × *LINEARE*. (*E. pseudolineare*, Hausskn.)

Specimens collected by Prof. Macoun at Ellis Bay, Anticosti, have been referred here by Dr. Trelease.

(11.) *E. DAVURICUM*, Fisch.

"Habit of *E. palustre*; stems terete or with occasional low decurrent lines; seeds fusiform, prominently beaked."

A span or two high mostly simple, the very slender stem sparingly incurved-pubescent, otherwise glabrous; roots densely fascicled; leaves less than 15 mm. long, somewhat crowded at base, alternate and remote above, linear or oblong, obtuse, remotely denticulate, sessile 1-nerved; flowers pale not very numerous, nodding; capsules erect