

Weeds, with their chief characters—*Continued.*

Colour, Size, Arrangement of Flowers and other Characters of the Plant.	Method of Propagation and Distribution.	Place of Growth, and Products Injured.	Methods of Eradication.
Purple, $\frac{1}{2}$ -in., umbel; leaves feathery pinnatifid; styles in fruit elongated, twisting spirally when ripe, bearded inside.	Seeds.....	Gardens, fields....	Hoe, cultivate....
Purple, $\frac{1}{2}$ -in., solitary; ripe pods black; seeds round, $\frac{1}{2}$ -in., mottled or velvety black.	Seeds, in grain	Fields ..	Summer fallow, hoed crop.
Violet and blue, $\frac{1}{2}$ -in., spikes long crowded 1-sided, about 30 flowers; pods, light brown; seeds round, $\frac{1}{2}$ -in., dark brown, mottled.	" in hay..	Hay fields, fields..	Plough, hoed crop.
Whitish, $\frac{1}{2}$ -in., spikes peduncled; pods oblong, $\frac{1}{2}$ -in., covered with hooked prickles.	Seeds, pods attached to stock, in hay	Summer fallows, pastures; wool.	Summer fallow early.
Rose coloured, small, in dense terminal panicles; leaves below brownish and woolly.	Seeds	Mountain pastures	Pull and grub out.
Yellow, $\frac{1}{2}$ -in., leafy cymes; leaves 8-parted; whole plant dark green, hairy.	"	Summer fallows, grain fields.	Summer fallow, cultivate.
Yellow, $\frac{1}{2}$ -in., stems spreading, ascending, paniculate, many flowered; leaves dark green above, silvery white beneath.	"	Pastures, lawns, hay fields.	Breaksod, cultivate
Bright yellow, $\frac{1}{2}$ -in., solitary on long stalks.	" runners.	Low lands, particularly if alkaline.	Summer fallow, cultivate.
Pink to deep rose, $\frac{1}{2}$ -in., corymb...	" rootstocks.	Summer fallows, grain fields.	Summer fallow early, harrow, cultivate.
Purple, in close compound cymes, 2-3-in. across; whole plant fleshy.	Seeds, portions of stem or root.	Pastures, hay....	Spud, break sod, cultivate.
Purplish, $\frac{1}{2}$ -in., panicle erect; leaf stalks very short; stem glandular pubescent.	Seeds, wind...	Low land, summer fallows, grain on stubble.	Plough fall or spring, summer fallow early.
Yellow, $\frac{1}{2}$ -in., open at night, leafy spike.	"	Summer fallows...	Pull, plough fall and spring.
White, turning pink, 2-in., malodorous leafy spikes; buds nodding; stems glistening white, simple, branched at the top.	Seeds and rootstocks.	Grain fields on knolls.	Summer fallow, cultivate thoroughly.
White, in umbel, central flower red; umbel 3-in. across, closing in like a bird's nest when mature; seed bristly.	Seeds, carried by animals.	Fields, pastures, road sides.	Break sod, spud.
White; umbel 2-in. across; seeds oblong, ribbed, smooth, aromatic.	Seeds	Road sides, pastures.	Spud, mow in flower
White; umbel 4-in. across; stem stout, spotted with purple, strong smelling, very poisonous.	Seeds, carried by floods.	Wet meadows, troublesome in hay, poisonous to stock,	"