

These different names refer to the same species. The wings are scalloped, not entire, and I accept Dr. Packard's generic determination with hesitation.

*Tornos infumataria*, n. s.

♀. Larger and stouter than *robiginosaria*, with the wings more elongate. Entirely smoky blackish. Fore wings with two sub parallel, oblique, irregular, black median lines, the exterior partly lined on the outside with whitish, but very faintly so. Hind wings with an indistinct mesial line, which is seen to be scalloped in the best marked specimens; beneath without markings. This concolorous species differs from its ally by the course of the waved median lines on primaries. Expanse 30 mil. June 3, 5; Texas (Belfrage, No. 604).

*Aspilates pervaria* var. *interminaria* Grote.

♂ ♀. Both sexes of this form, which is smaller than the type and differs at once by the absence of the lines on primaries, have been collected by Mr. Belfrage in Texas (male, June 5, No. 602; female, May 22, No. 653). It is paler than the type and looks like a different species.

---

LIST OF BOMBYCIDÆ OCCURRING ON THE ISLAND OF  
MONTREAL, P. Q.

BY F. B. CAULFEILD AND C. W. PEARSON, MONTREAL, P. Q.

---

BOMBYCIDÆ.

Lithosiinæ.

*Hypoprepia fucosa* Hübn. Not common.

*Euphanessa mendica* Pack. Common; end of June, July.

*Crocota Treatii* Grote. Very rare; C. W. P.

“ *aurantiaca* Hübn. Very rare; C. W. P.

“ *brevicornis* Walk. Rare.

*Utetheisa bella* Linn. Rare.

Arctiinæ.

*Callimorpha Lecontei* Bois. Common; end of June, July.