

-
204. *Estola sordida* LeConte.
Found at San José del Cabo, Lower California.
205. *Peritapnia nudicornis* Bates.
Found in Sierra El Chinche, Lower California.
206. *Lysimena californica* Horn.
Found by Van Dyke at Santa Monica, by Fall rarely at Pasadena, and by Fuchs in Yuba County.
207. *Pogonocherus crinitus* LeConte.
Bred from oak at Pasadena by Dr. Fenyes; bred from oak in Marin and Alameda Counties by Van Dyke. Found in Middle and Southern California. June 2–August 20.
208. *Pogonocherus concolor* Schaeffer.
Beaten by Pilate from *Fremontidendron*. Taken by Schaeffer in the middle Sierras.
209. *Pogonocherus oregonus* LeConte.
Found in the middle Sierras. Taken by Nunenmacher in Eldorado County; taken by Fall at Tahoe; thought by Van Dyke to breed in fir, probably also Douglas spruce. May 10–August 9.
210. *Pogonocherus californicus* Schaeffer.
Found in the middle Sierras; taken by Van Dyke in Placer County; taken by Blaisdell in Calaveras County; taken by Schaeffer in Tulare County. June 6.
211. *Pogonocherus volitans* LeConte.
Found at Cape San Lucas Lower California.
212. *Ataxia setulosa* Fall.
Found in Lower California.
213. *Saperda hornii* Joutel.
Taken at Los Angeles, Northern California, and the Sierras. Taken from willow by Van Dyke and Rivers.
214. *Saperda populnea* Linné.
Found in Northern California and as far south as Los Angeles. Feeds on poplar. Taken by author by beating poison oak bushes at Felton, Santa Cruz County. July 1–4.
var. *tulari* Felt & Joutel.
Taken in Merced and Tulare Counties.
215. *Mecas inornata* Say.
Breeds in the stems of the false sunflower and in *Helianthus tuberosus*.