

recalls the dry earth system regarding which great expectations were at one time entertained. The advantages of moss litter over dry earth for the purposes in question are, however, very decided. They consist in the perfect inoffensiveness of the moss litter product, in the fact that one part of moss litter will deodorize and dry at least six parts of mixed excreta, and in the greater agricultural value of the resulting manure. Dry earth (which is required in quantity at least equal to that of the excreta) is valueless from an agricultural point of view, but this is not the case with moss litter, which as its analyses show, often contains as much nitrogen as ordinary barn-yard manure. Numerous analyses have been made of moss litter manure as produced in Germany, and its average contents from seven different towns may here be stated.

	p. cent.	lbs. per ton.		Value per ton.
Nitrogen.....	0.664	13.28	at 13c.	\$1 72
Phosphoric acid.....	0.350	7.00	5	0 35
Potash.....	0.285	5.70	5½	0 30
Water.....	83.00			\$2 37

Numerous trials have been made on various crops with this manure, and very satisfactory results are always reported. In all cases it is stated to excel barn-yard manure even when the latter is used in much greater quantity.

Canada possesses in its bogs and swamps inexhaustible quantities of moss litter which is frequently found in beds, several feet in thickness lying above the peat. The following tests have been made in the Inland Revenue Laboratory, of moss litter from various localities in the Dominion:—

	Moisture.	Ash.	Nitrogen.
	p. c.	p. c.	p. c.
Moss litter, Berwick, N.S.....	14.40	1.16	1.26
Black muck, Berwick, N.S.....	13.30	3.68	1.58
Moss from Great Village, N.S.....	63.44	3.46	0.63
Sphagnum moss from Shippegan, N.S.....	12.45	1.55	0.55
Light coloured moss litter from Lincoln Parish, N.B.....	11.55	1.40	1.79
Dark coloured sample from the foregoing locality.....	10.95	0.80	1.06
Moss litter from Musquash, N.B.....	11.50	0.95	0.82
Moss litter from lower layer.....	12.50	0.90	0.72
Peat from St. Bridget, P.Q.....	13.30	2.50	1.48
Peat from St. Hubert, P.Q.....	12.35	2.68	1.84
Light coloured moss litter from Caledonia Springs, Ont.....	10.00	1.60	2.95
Dark coloured moss litter from the same locality.....	11.60	2.70	2.23
Peat from the same locality.....	10.95	3.80	2.94
Surface moss from the Mer Bleu at Eastman's.....	10.85	2.80	0.71
Surface moss from the Mer Bleu at Baldwin's Farm.....	7.90	2.65	1.47
Surface moss from the Mer Bleu at Baldwin's Farm 18 inches deep.....	27.90	1.72	1.64
Peat from Mer Bleu at McFadden's Farm, Navan, wide ditch.....	22.60	4.40	2.21
Peat from Mer Bleu at McFadden's Farm, Navan, narrow ditch.....	9.40	6.62	2.80
Peat from near Stratford, Ont.....	16.80	9.10	1.91
Hypnum moss from the Ellice bog, Stratford, Ont.....	8.75	9.72	2.01
Moss litter from Welland marsh, Ont.....	3.85	4.70	1.51
Peat lying underneath the foregoing.....	5.30	4.83	1.41
Peat from same locality, 4½ feet deep.....	3.25	41.25	1.52
Peat from Dobson's bog near Beaverton, Ont.....	18.42	9.04	1.89

The manufacture of moss litter has been attempted at Musquash, in New Brunswick, and also in Welland County, Ontario. From the latter locality I was supplied with several bales of the moss litter for experimental purposes, and Dr. LaBerge, of Montreal, undertook to superintend the carrying out of an experiment to determine its deodorizing and absorbent qualities. He reported that 100 pounds of moss litter were