

Bulimina pupoides D'Orb., rare.
Bolivina levigata (Will.), very rare.
Cristellaria Italica, Defr. (young), very rare.
Truncatulina Sp., very rare.
Rotalia orbicularis, D'Orb., very rare.
 Radiolaria frequent.
 Sponge spicules, rare.

No. 3. Boulder clay near Victoria. Weight 3 lbs. 1.6 oz. Troy. After washing—Fine 1 lb. 0.1 oz. Coarse 0.6 oz. Stones mostly rounded, some angular.

Gaudryina Sp., frequent.
Bulimina pupoides, D'Orb., rare.
Rotalia orbicularis, D'Orb., frequent.
Nonionina depressula (W. & J.), very rare.
 " *scapha* (F. & M.)? very rare.
 Radiolaria, frequent.
 Sponge spicules, rare.
 Ostracod, very rare.

It may be of interest to add for comparison, the species actually recognized by Mr. Wright in the several small samples of Cretaceous material supplied to him. These are, nearly in order of relative abundance, as follows:—

Textularia globulosa Ehr., very common.
Globigerina Cretacea, D'Orb., very common.
 " *digitata*, Brady, rare.
Anomolina ammonoides, (Rss.), rare.
Nodosaria Zippelii, Rss., very rare.

GEORGE M. DAWSON.

GEOLOGICAL SURVEY OF CANADA,
 March 10, 1897.