


DR. A. C. DANIELS

WORLD FAMOUS

COLIC CURE



You can cure any case of Colic if you use
this celebrated remedy in time

It is a fact that twenty-seven horses die of colic where one is burned to death.

You insure against fire. Do you insure against colic? Dr. Daniels' Colic Drops are the one safe policy to hold against Horse Colic. It is always quick to act, safe and convenient to administer, sure to save the horse when given

promptly; a woman or a child can save a horse, so simple is its use.

Symptoms of Spasmodic Colic.—Distended Bowels, relieved by pressure; pulse natural; lies down and rolls, and seems easier at intervals; legs and ears generally warm; comes on suddenly; is never preceded, and rarely accompanied by fever; looks back at flanks; paws and kicks; will not drink.



A Symptom of Colic

Symptoms of Flatulent Colic.—Similar to those of spasmodic, except the PAIN IS CONTINUOUS, not so violent; abdomen distended VERY HARD with drum-like sound when struck, followed by difficult breathing; perspiration; trembling of limbs; sighing respiration; pulse rapid; staggers from side to side; bowels constipated; lies down with caution.

Horse Colic is Caused

By a great variety of conditions. The most common are indigestion, change of feed or water, drinking cold water on an empty stomach, when tired or exhausted, constipation, feeding new hay, oats or corn, any grain that has soured, ulcers, worms, etc. Many cases are brought on by the horse taking a sudden chill by standing in a draft when heated; also by over-feeding, which causes an unusual amount of gas to accumulate; neglect of the usual evacuations, such as going a long time without urinating. All of the above, as well as many other causes, or a combination of them, will bring on colic.

Treatment

Our treatment will seem so easy and simple, as compared with former practice, as to raise some doubts of its results in the minds of people who have had sad experience with other medicines; but follow directions faithfully and the result will please you. Get the horse into a nice spot of soft turf or into a deeply bedded box stall, so that if he throws himself down violently he will not bruise his flesh or rupture himself. Give as soon as possible a dose out of No. 1 bottle, 30 drops (the syringe full), as far back on the tongue as possible. In ten minutes give the same sized dose out of No. 2 bottle, and then continue giving these doses as above, first one and then the other, about ten minutes apart. If a horse is very sick before you