

2. *Microdon viridis*, Townsend.

Ainsworth, July 11, 1903, by R. P. Currie, and Vancouver, June 11, 1904, by R. V. Harvey.

3. *Microdon marmoratus*, Bigot.

A single specimen taken at Vernon, Aug. 14, 1904, by R. V. Harvey.

4. *Chrysotoxum ypsilon*, Williston.

Kaslo, June 5, 1903, by H. G. Dyar. (Identification by Coquillett.)

5. *Pipiza pisticioides*, Williston.

Taken at Kaslo by Mr. Currie on various occasions ranging from June 12 to July 20, and by Mr. Cockle, June 30, 1906.

6. *Pipiza nigripilosa*, Williston.

A single male specimen taken at Vancouver, June 21, 1904, by Harvey, differs from eastern specimens in having the eyes dark pilose instead of light pilose, but otherwise the agreement is very close.

7. *Pipiza quadrimaculata*, Panzer.

Five specimens taken at Vancouver, July 2, 1906, by R. V. Harvey and R. S. Sherman. This European species has not been recorded from North America until very recently. In "Psyche" for August, 1907, Mr. C. W. Johnson notes its capture at North Mountain, Penn., and Mt. Greylock, Mass. The species thus evidently furnishes another example of circumpolar distribution to be added to the already long list among the Syrphidæ. It is easily distinguished from other American species of the genus by the two interrupted yellow cross-bands on the abdomen, forming the four spots which suggest the specific name.

8. *Chilosia chalybescens*, Williston.

Grouse Mt., July 1, 1904, one specimen by R. V. Harvey.

9. *Chilosia occidentalis*, Williston.

Two specimens by R. V. Harvey, one from the Hope Mts., July 24, 1906, the other at Vancouver, June 22, 1906.

10. *Chilosia hoodiana*, Bigot.

One female specimen from Similkameen, July 20, 1906, by R. V. Harvey, I place here, though it shows some differences. The yellow of the knees ("geniculis tibiærumque basi, fulvis," Bigot) is almost wanting, and the fourth abdominal segment is entirely shining.