

*Nysius strigosus*, Uhler.

= *senecionis*, Baker (nec Schilling).

*Cymus discors*, Horváth (n. sp.).

= *tabidus*, Provancher.

= *claviculus* of American authors, not of Fallén.

*Cymodema exiguum*, Horváth.

= *tabidum* of American authors, not of Spinola.

*Ischnorhynchus geminatus*, Say.

= *Cymus franciscanus*, Stål.

= *Ischnorhynchus didymus* of American authors, not of Zetterstedt.

= *I. resedæ* of American authors, not of Panzer.

*Aphanus umbrosus*, Distant.

= *Dorachosa illuminata*, var. *umbrosis*, Dist.

= *Microtoma carbonaria*, Uhler (not Rossi).

= *M. atrata* of American authors, not of Goeze.

*Uhleriola*, Horváth, new genus to contain *Rhyparochromus floralis*, Uhler, which does not pertain to that genus, nor even to the division *Aphanaria*.

*Emblethis vicarius*, Horváth (n. sp.).

= *arenarius* of American authors, not of Fieber.

= *griseus* of American authors, not of Wolff.

= *Gonianotus marginepunctatus*, Uhler (not Wolff).

*Melanorhopala clavata*, Stål.

= *Cantacader Henshawii*, Ashm.

The Ashmeadian types were seen by Dr. Horváth when in Washington. This is sometimes wrongly attributed to the genus *Lasiacantha*, following Lethierry and Severin (Cat. Gen. Hém., III : 18).

*Aradus crenatus*, Say.

= *dilatatus*, Dufour.

*Hypselosoma*, Reuter.

= *Glyptocombus*, Heid.

*Orthotylus chlorionis*, Say.

= *Phytocoris flavosparisus*, F. Sahlberg.

*Acanthia xanthochila*, Fieber.

var. *limbosa*, Horn. A European form, now recorded for the first time from America.

*Empoasca aspersa*, Gillette & Baker.

= *tesselata*, Gillette (not Lethierry).