


I will call, out of charity to myself, "ill-luck" as I suffered in connection with this patient; they all arose out of a bad commencement. In giving the patient a list of two or three lodging-houses, I included one which, though excellent in itself, was not a good one for our particular purpose. I recollected this immediately; but, expecting to see the patient next day, and having arranged not to operate for a day or two, I thought I should have an opportunity of setting things right. Unfortunately, I did not hear again from my patient till the eve of the date fixed for the operation, and he had in the meantime taken rooms with an exceedingly bad light. A corner window on the ground floor, looking into the bottom of a kind of pit, and the enormous bed in the room, when placed in such light as there was, blocked the way in such a manner as almost to paralyze the nurse and greatly interfere with the assistant. Further, the eventful morning was dark and dull, even for London; and we had to use candle light. My assistant was more than fully occupied in endeavors to retract the fatty walls of the wound; these were so thick and deep that it was not until I had made an incision three if not four times as long as the usual one, that I could get proper access to the inguinal ring and its pillars. When I now proceeded to put in the thick catgut suture with a handled needle, the shank of the latter bent with the effort necessary to bring the eye into view after it had passed through the pillars. There was a good deal of oozing, and the nurse being occupied in holding the candle, while the assistant's hands were monopolised by the retractors, I had to sponge for myself. When the suture was properly inserted, I paused to reflect ere I injected a powerful irritant into the inguinal canal, after all the parts had necessarily been subjected to much rough usage.

Had we been engaged on some necessary operation, such as an amputation, to go on and make a thorough finish of it would have been a matter of course; but here was a very different state of things. My main object in doing these operations of injection is to find a really safe and reasonably certain mode of doing the radical cure of hernia. I would therefore infinitely rather fail to do any good than risk doing any harm. Therefore, considering the unusual stoutness of my patient, the large wound it had been necessary to make, the rough usage to which it had been subjected, the unreliableness of the antiseptic precautions which it had been possible to take under the circumstances, and the doubt whether, even under the most favorable conditions, injection would do much good to a hernia coming through such an immense aperture (it would admit four fingers), I determined to refrain from exposing my patient to the risks associated with injection into his canal. Had I had him on the hospital operating-table, in an excellent light and surrounded by plenty of assistance,

I should probably have there and then tied the neck of the sac, excised the fundus of it, sutured the ring, and confidently expected a good result. But this course was here at present out of the question, although I had obtained my patient's leave to do as I liked. I therefore merely put a little of the injection on the sutured ring, placed a drainage-tube, sponged and cleansed the wound with 10% sublimate lotion, and dressed with turf-moss dressing, etc., sublimated in the usual way.

Very glad I was afterwards that I had refrained from putting the injection into the canal, otherwise I should have probably had to deal with an abscess in the abdominal walls between the muscles, for the wound did not heal by first intention, and there was a good deal of suppurating with some sloughing. In addition to this, an attack of pneumonia with rusty sputa, high temperature, etc., developed in a few days, and kept the patient incessantly coughing, besides causing me some



anxiety, and compelling me to pay the most rigorous attention to the wound, with a view to preventing septic absorption. I used to dress it three times a day. The pneumonia was partly aggravated by his being kept always on his back with the foot of the bed raised, and disappeared rapidly as soon as I allowed him to sit up a little. The hernia remained up for a fortnight, and then came down in a fit of coughing.

The suture had given way, and I removed it through the wounds, which was still unhealed. I then let him sit up in bed, and all signs of the pneumonia quickly vanished, showing that whatever had been the original cause, it had been kept up and increased by the statical effect of the continued supine position with the lower extremities