

except a small wedge in front, post-scutellum, a large irregular diagonal spot on mesopleura, spot beneath posterior wing, circular spot just back and above, semicircle around insertion of abdomen with a ray extending forward from each coxa, and a third from the abdomen, base and apex of the first two abdominal segments, 4 anterior coxæ and trochanters, stripe on posterior coxæ above and second joint of posterior trochanter, yellowish-white. Posterior coxæ and first joint of trochanter black; posterior tarsi dusky at tips; remainder of legs honey-yellow. Antennæ black, fulvous toward the tip. Wings hyaline, infumated at the apex; areolet small, petiolate. Entire thorax coarsely, but not closely, punctured. Abdomen smooth, shining.

Described from 2 ♀♀ collected at the Michigan Agricultural College.

*MENISCUS MICHIGANENSIS*, n. sp.

♀. Length, 10 mm.; ovipositor, 8 mm. Black, with yellow markings; abdomen beyond second segment rufous. Broad orbital lines, entire front, scape beneath, annulus on antennæ, mouth except tips of mandibles, gula, entire prothorax, tegulæ, line beneath, large triangular spot in front confluent with a cuneiform line on mesonotum, extending back nearly to the scutellum, venter except anterior part of mesosternum, with a continuation of the same on to the pleuræ, ending in a large hook beneath the wings, metapleuræ and two lines above joining posteriorly, base, apex and a large spot in the centre of the first abdominal segment, base and apex of second segment, connected by a longitudinal line, lemon-yellow. Posterior femora rufous; 4 posterior trochanters black at base, and coxæ with black stripe above; 2 hind coxæ black inside; extremity of posterior tibiæ, and basal joint of tarsi, except tip, black; remainder of legs lemon-yellow. Wings hyaline without areolet. Antennæ brown, black above. Thorax and face coarsely, but not closely, punctured. Abdomen smooth, shining.

Described from 1 ♀ taken at the Michigan Agricultural College.

? *MENISCUS JOHNSONII*, n. sp.

♀. Length, 11 mm.; ovipositor, 1.5 mm. Black, with yellowish-white markings, as follows: Entire head, except occiput, spot enclosing ocelli and antennæ, narrow central line beneath, reaching to clypeus, and tips of mandibles; prosternum; mesothorax with two cuneiform lines on margins of mesonotum, two stripes in front of scutellum, scutellum except posterior part and spot in front, two spots beneath primary wings and