

profession have appreciated its worth and thorough reliability is well apparent, from the place it holds in the regard of every physician who appreciates stability and honesty.—*The International Journal of Surgery*.

---

SANMETTO IN ENLARGED PROSTATE AND CHRONIC CYSTITIS, IRRITABLE BLADDER AND URETHRA.—I have used Sanmetto in enlarged prostate and chronic cystitis in old men, with marked good results, and observed that there was decided aphrodisiac effects; also in irritable bladder and urethra in the early months of pregnancy, with very happy results.—*M. A. Rush, M.D., Anderson, Ind.*

---

FERMENTATIVE DYSPEPSIA.—I have been treating a case of Fermentative Dyspepsia for some time, and thought I would wait until the cure was certain and likely to be permanent before I wrote you, but as it is now four weeks since patient has taken anything in the shape of medicine and has had no trouble whatever during that time (although rather careless about his diet), I thought it would be safe to inform you of one more victory for Glyco-Thymoline.

The patient, age 60, occupation—farmer. As far back as he can remember has had trouble with his stomach; heartburn, sour stomach, and enormous collections of gas in stomach and bowels. This is his own way of expressing his troubles. His condition when I first saw him was about as we described it above. The collections of gas in stomach and bowels were so enormous that he could scarcely ever lie down. Spent his nights in a sitting position. He was very much emaciated and completely discouraged, did not consider life worth living, and, indeed, had made several threats of suicide.

Well, it was not a very promising case for any physician to take, but in thinking it over I made up my mind that this would be a pretty good opportunity to try Glyco-Thymoline internally, and so I agreed to do what I could for him. To begin with, I regulated his diet, and then, as he was very constipated, got his bowels to acting regularly, one evacuation every day. I then, every other day, washed out his stomach with one ounce of Glyco-Thymoline in a pint of tepid water and gave him a teaspoonful of Glyco-Thymoline in a tablespoonful of water before and after each meal, and directed him if he was much troubled with gas in the night to take another dose about midnight.