



VOL. XXIII.

MONTREAL, NOVEMBER, 1894.

edical

No. 2.

ORIGINAL COMMUNICATIONS.

| Three Cases of Cœliotomy : An Ova- rian Cyst; A Tubal Pregnancy : and a Hematoma of the Ovary Double Femoral Herniotomy in a Woman 64 years of age; Primary Union of Both Wounds | 25 28 |
|---|-------------|
| Extract of President's Address be- | |
| | 28 |
| fore Nova Scotia Medical Society . | 28 |
| | |
| SOC)ETY PROCEEDINGS. | |
| The Montreal Medico-Chirurgical | |
| Society | 31 |
| A Case of Symphysiotomy | 31 |
| Old Dislocation of the Hip-Joint | · · |
| Treated by Resection | 31 |
| Resection of the Intestines | 32 |
| A Case of Pylerectomy | $\tilde{2}$ |
| An Appendix containing an Ordinary | _ |
| | |
| Pin as the Exciting Cause of Per- | |
| forating Appendicitis | 33 |
| Calcareous Tumor of the Thyroid | |
| producing Esophageal Obstruc- | |
| Tion | 33 |
| Tumor of the Prostate | 34 |
| Eaur Calculi weighing 5 oppose 1 | |

| Four Calcul | weighing | 5 ounces 1 the Bladder | |
|-------------|------------|---------------------------|---|
| drachm rer | noved from | the Bladder | 3 |

Griginal Communications.

- *THREE CASES OF CŒLIOTOMY ; AN OVARIAN CYST ; A TUBAL PREGNANCY ; AND A HŒMA-TOMA OF THE OVARY.
- By A. LAPTHORN SMITH, B.A., M.D., M.R.C.S. ENGLAND, F.O.S. LONDON, Fellow of the American 'Gynæcological Society, Gynæcologist to the Montreal Dispensary, Surgeon to the Women's. Hospital.

CASE I. Multilocular cyst of right ovary. Removal. Recovery.

This tumor, which measures about 15 inches in length by about 9 in thickness, in its present dried condition was very much larger before its removal. The patient, Mrs. T., from whom it was removed, was sent to me from Valleyfield, on the 4th of August, the diagnosis having been made by her physician. It had a distinct ovarian expression. An

* Read before the Medico-Chirurgical Society, Montreal, 2nd November, 1894.

CONTENTS.

Romen Medical Society 3 PROGRESS OF SCIENCE.

| 100000000000000000000000000000000000000 | |
|---|----------------------|
| Cancer a Local Disease New Observations in Gonorrhoa | 38 38 |
| Anæsthesiä Growing Pains | 38 38 |
| Treatment of Hydrocele | -38 |
| Accentuation of the Pulmonary | 38 |
| Sound in Perityphlitis Rare Case of Traumatic Cyst of the | -58 |
| Stomach | 39 |
| Homes for Consumptives New Method of Examining for Renal | 39 |
| Calculus | 40 |
| Primary Cancer of the Gall-bladder. The Urine in Eateric Fever | 40 40 |
| An Extreme Case of Ascites | 40 |
| Renal Hæmorrhage from an unusual Source | 40 |
| Water in Typhoid Fever | 41 |
| Treatment of Red Nose Second-Hand Syringes | 41 -41 |
| Kentucky School of Medicine | 42 |
| Dangers of the Long Rectal Tube. The Best Treatment of Hemorrhoids | - 4 2 - 42 |
| THE POST THREE OF CLENTING | |

EDITORIAL.

| County Societies | 43 |
|---|----|
| The Death of Oliver Wendell Holmes | 45 |
| American Public Health Association The American Electro-Therapeutic | 45 |
| Association | 46 |
| | |
| BOOK NOTICES. | |
| The Year-Book of Treatment for 1894 | 47 |
| The Retrospect of Practical Medi- | |
| cine and Surgery | 47 |
| Pamphlets received | 47 |
| and the second se | |
| Class-Room Notes | 48 |
| Publishers' Department | 48 |

ovarian tumor is often mistaken for pregnancy; but in this case, the patient was led to adopt this view of her case, more especially because there was a hard, oval lump in the right hypochondrium which, I must admit, felt very like a child's head. She had begun to menstruate at 15, nor-. mally, was married at 21, and had been married two years without having become pregnant, although she had skipped a period twice since her marriage. She had first noticed her abdomen enlarging sixteen months ago. On examination, all the evidences of an ovarian cyst were found, and the uterus, which was pushed backwards and to the left, measured three inches in depth. She entered my private hospital on the 1st of September, and the tumor was removed on the 4th, without any difficulty. The wound healed by first intention, and she made such a rapid and easy recovery, that she was out of bed in two weeks and went out in twenty-five days. The other ovary was healthy, and was not removed. CASE II. Tubal Pregnancy. Removal. Recovery. The patient from whom this specimen was removed was Mrs. W., æt.