

rarely with two large yellowish spots on each.

.14-.15 in. *quercata*, Fabr.

While *flavocyanea* is included in the above table, on account of its being recorded in the Society's list, it has probably been identified in error, since it is a Southern species.

DISONYCHA, Chevr.

Also contains large or moderate sized species, some of them even exceeding *Edionychis*, which they often resemble in markings, but they may easily be separated therefrom by the claw-joint of the hind tarsi not being swollen. They separate thus:

A. Elytra not striped.

Thorax yellow with three black spots arranged in the form of a triangle; under surface of body and the legs black .20-.25 in. *triangularis*, Say.

Thorax yellow, not spotted; abdomen yellow, femora usually yellow at basal half. .21-.23 in. *xanthomelæna*, Dalm.

AA. Elytra striped.

b. Form very elongate; elytra vaguely grooved; thorax somewhat uneven.

Body beneath black, except sides of thorax, which are margined with yellow. Black spot on disk of thorax very large. var. *limbicollis*, Lec.

Body beneath partly black, abdomen paler at sides and apex, thorax with under surface entirely yellow, discal spot on upper surface smaller. .26-.30 in. *pennsylvanica*, Ill.

bb. Form not very elongate; elytra and thorax even, the former with discal and submarginal vittæ.

c. Abdomen densely punctured, conspicuously pubescent.

d. Head coarsely punctured from side to side. .22-.36 in. *quinquevittata*, Say.

dd. Head smooth at middle.

Elytral vittæ rather broad, head and body beneath more or less clouded with darker, labrum piceous. .22-.26 in. *crenicollis*, Say.

Elytral vittæ narrow, head and body beneath always pale yellow, labrum pale. .20-.26 in. *caroliniana*, Fabr.