veins pale, brachial vein towards its apex and a longitudinal vitta within the apex of the areole fuscous. Male and female. Length 7, width 2 mm. Klauck's four specimens taken with the preceding. This is the species found at Island Lake last year by Mr. Palmer, and listed by me as Neoborus, sp. I took it at Lake Placid, and have seen others from Montreal taken by Mr. Beaulieu.

 ${\it Poecilocapsus\ lineatus}, {\it Fabr.} {\it ---} Apparently\ common\ at\ all\ places\ along\ the\ Ottawa\ River.$ 

Camptobrochis grandis, Uhler.—Temagami, Temiskamingue and Nigger Point on Quinze Lake.

Lygus pratensis, Linn.—Abundant everywhere, with its variety flavonotatus, Prov.

Lygus invitus, Say. - Klauck's, on Quinze Lake.

Lygus monachus, Uhler.—One example from Barrier River.

Lygus pabulinus, Linn.—White Rapids, one example.

Adelphocoris rapidus, Say.—Apparently abundant at all stations. Dr. Reuter has separated this species and its allies from Calectris.

Neurocolpus nubilus, Say.—Temagami, one example.

Compsocerocoris annulicornis, Reut.—Barrier River.

Phytocoris pallidicornis, Reut.—Taken in numbers at all stations.

Phytocoris eximus, Reut.—Bear Island, Lake Temagami.

Lopidea marginata, Uhler.—This species seems to have been common, as Mr. Palmer brought home numbers from Temagami, Barrier River and White Rapids.

Stiphrosoma stygica, Say.—Temagami, Barrier River and Temiskamingue. These agree with material from the mountains of Colorado in being a little smaller than those taken about Buffalo.

Dicyphus agițis, Uhler.—Temiskamingue and Nigger Point, Quinze Lake.

Hyaliodes vitripennis, Say.—Mr. Palmer brought home from Barrier River one example of the dark form that, about Buffalo, is found on oak.

Cyrtorrhinus marginatus, Uhler.-Taken at Barrier River.

Orthotylus, sp.—Quinze Lake, one example.

Oncotylus chlorionis, Say.—Temiskamingue, one example. Plagiognathus obscurus, Uhler.—Abundant at all stations.

Plagiognathus sp.—One example of a small species, with a vittate scutellum, was taken at Temiskamingue.

GERRIDÆ.

Gerris remigis, Say.—One brachypterous example was taken at Quinze Lake.