

the ocellar furrow deep, the interocellar furrow deep and Y-shaped; the saw-guides convex above and below, slightly, obliquely truncate at apex, with rounded angles; body black, with the collar, tegulae, front and middle legs below the knees, and the base of the hind tibiae, white. Length, 7 mm.

Habitat.—Colorado. Professor C. F. Baker, collector.

*Empria contorta*, n. sp.—Female. Clypeus with a median carina and without a median tooth, shallowly emarginate, lobes broadly rounded; antennal furrows interrupted; ocellar basin broad, oval, with sloping sides on its lower three-fourths, constricted just below the median ocellus, and expanding slightly at the ocellus; the median fovea broad and deep, located near the middle of the lower portion of the ocellar basin; the third segment of the antennae longer than the fourth; the ocellar and interocellar furrows deep, the interocellar furrow somewhat Y-shaped; the saw-guides strongly concave above, slightly convex below, broadly rounded at apex, with a broadly rounded knob at apex above; body black, with the collar, tegulae and legs below the knees white. Length, 7 mm.

Habitat.—Chimney Gulch, Colorado. Mr. E. J. Osler, collector.

*Empria culpata*, n. sp.—Female. Clypeus with a median carina, shallowly, roundly emarginate, without a tooth, lobes rounded; ocellar basin extending from the median ocellus to the dorsal margin of the antennal foveae, a linear depression, broadest below; the median fovea shallow, quadrangular, separated from the ocellar basin; ocellar and interocellar furrows distinct; the third segment of the antennae longer than the fourth; the saw-guides straight above, convex below, obliquely rounded at apex, densely hairy; body black, the collar white, the legs below the knees more or less rufous infuscated. Length, 6 mm.

Habitat.—Olympia, Washington. Professor Trevor Kincaid, collector.

*Empria cumulata*, n. sp.—Clypeus with a median carina, roundly emarginate without a tooth, lobes rounded; the ocellar basin a narrow linear depression extending from below the dorsal margin of the antennal foveae to the median ocellus, somewhat interrupted at middle, the median fovea deep, elongate; the ocellar and interocellar furrows indicated; the postocellar area polished; the third segment of the antennae slightly longer than the fourth; the saw-guides straight above, convex below, obliquely truncated at apex; body black, with the collar white, the front and middle legs reddish infuscated below the knees. Length, 6 mm.

Habitat.—Olympia, Washington. Professor Trevor Kincaid, collector.