

by blood-streaked sputum. On March 28th patient went out of doors, but the snow caused eyes to ache and headache. This illness was diagnosed measles.

March 31st to April 2nd.—Continued poorly. Vomited a half basinful of straw-colored fluid. His headache was relieved by vomiting. Vomited three times while dressing on April 1st. It was very difficult to stoop down to lace his shoes. When dressed he felt drowsy and had severe frontal headache. Felt weak in the legs and felt like falling forward, but not to one side or the other. After a little time was able to walk. Came to Toronto April 2nd, went to bed at 7 p.m., and vomited about a quart of coffee-like watery fluid. The attacks of vomiting came on suddenly with a good deal of retching, but the vomitus was not projected. Felt feverish and had a good deal of frontal headache. Felt better next day, but had slight headache.

Admitted to Toronto General Hospital.

Condition since admission:—

Has had headaches ever since, which came on after lying down, and which became less intense by rising and walking about the ward. These headaches felt as though there was a heavy weight inside trying to shove its way out and situated in the low frontal region and to some extent on both sides. Has not vomited since admission. The pain in the back has persisted ever since the attack of supposed grip. He does not notice it when walking, only when bending down or getting up from a chair. The blowing sound in the left ear has persisted ever since the attack of supposed grip and is more pronounced when he is lying awake at night.

PHYSICAL EXAMINATION.

On percussion of the head most pain is felt over an area the size of a fifty cent piece to the left of the median line and at the level of a line drawn upward, perpendicular to the zygoma, one inch in front of the external auditory meatus. The percussion note is higher pitched and shorter. Face and head are well formed, and two sides fairly symmetrical.

Nervous system.—Intelligence good; not emotional or irritable. No delusions nor hallucinations. Mental condition has not changed in the last year. Can read for about five minutes, after which time things become so dim that he can no longer read. When in the Home in England, while walking in his sleep, he fell, cutting his forehead. There is no paralysis of any group of muscles. There is rather a coarse fibrillar tremor of the tongue and of the extended fingers. There is no facial tremor when the muscles are tightly