

LILIACEAE.

Lloydia serotina (L.) Reieh.

Camden bay—Herschel island.

COLCHICACEAE.

Tofieldia palustris Huds. [*T. borealis* Wahlenb.]

Bernard harbour.

DICOTYLEDONEAE

SALICACEAE.

Salix anglorum Cham.

Cape Bathurst—Young point—Bernard harbour—Wollaston land (F. J. and D. Jenness)—Sandstone rapid, Coppermine river—Cape Barrow and Bathurst inlet (R. M. Anderson).

S. reticulata L.

Camden bay—Collinson point—Herschel island—Young point—Bernard harbour—Epworth harbour (J. R. Cox and J. J. O'Neill)—Wollaston land (D. Jenness)—Cape Barrow (R. M. Anderson)—Taylor island, Victoria strait (J. F. Bernard).

S. fullertonensis C. Schn.

Kater point and Arctic sound, Bathurst inlet (R. M. Anderson).

S. niphoclada Rydb.

Icy reef.

S. ovalifolia Trautv.

West of Martin point.

S. ovalifolia Trautv. var. **camdensis** C. Schn.

Camden bay—Collinson point—West of Martin point.

S. phlebophylla And.

Sadlerochit river.

S. pulchra Cham.

Camden bay—Collinson point—Herschel island—Bernard harbour.

S. rotundifolia Trautv.

Camden bay—Collinson point.

S. Richardsonii Hook.

Camden bay—Sadlerochit river—Herschell island—Croker river, Dolphin and Union strait (R. M. Anderson)—Clay banks below Bloody falls on east side of Coppermine river—Wollaston land, on sand flats.

Populus tremuloides Michx.

Arctic Alaska. Hulahula river—Near Sandstone rapid, Coppermine river, (R. M. Anderson).

BETULACEAE.

Betula glandulosa Michx.

Demareation point—Herschel island—Bernard harbour—Lady Franklin point, Wollaston land.