

- Tonsillitis, Follicular (continued)**  
Peroxide of hydrogen solution in, II. 21
- Tonsils, Diseases of.** See **Rheumatism**, Chronic muscular, II. 460—64
- Torticollis.** See **Rheumatism**, Chronic muscular, II. 460—64
- Tracheotomy** for diphtheria, II. 595—96  
for oedema of the glottis, I. 554
- Trichocerebellus dispar.** I. 305, 324
- Tubercular meningitis.** See **Meningitis, Tubercular,** II. 242—47
- Tubercular peritonitis.** See **Peritonitis, Tubercular,** I. 334—42
- Tuberculosis, Pulmonary.** See **Phtisis,** I. 715—II. 77
- Tumour, Intracranial,** II. 203—68  
Causation of, II. 265  
Focal symptoms, II. 266  
Forms of, II. 264  
Mercurial inunction in, II. 266, 267  
Potassium and sodium iodides in, II. 266  
Surgical measures in, II. 267  
Symptoms of, II. 266
- Typhilitis.** See **Perityphilitis,** I. 260, 261, 267
- Typhoid, or Enteric fever,** II. 600—706  
Acetone in, II. 688  
Alcohol in, II. 687  
Antiseptic treatment, II. 687  
Antityphoid serum in, II. 687  
Aperients in, II. 692  
Bacilli in, II. 669—72  
Bed-sores in, II. 705  
Boric acid in, II. 688  
Calcium lactate in, II. 708  
Calomel in, II. 688, 692  
Carbolic acid in, II. 688, 694  
Carbonate of guaiacol in, II. 688, 694  
Characteristic lesions of, II. 673  
Chlorine in, II. 688—90  
Chloroform in, II. 694  
Corrosive sublimate in, II. 680, 698  
Course of, II. 673—75  
Creasote in, II. 688  
Diarrhoea in, II. 701  
Diet in, II. 683—86  
Diffusion of, II. 671  
Haemorrhage in, II. 702  
High temperature in, II. 696—97  
Hydrotherapy in, II. 696—701  
Indications for treatment of, II. 682, 687  
Iodine in, II. 688  
Laparotomy in, II. 708
- Typhoid or Enteric fever (continued)**  
Maintenance of patient's strength in, II. 692  
Naphthalin in, II. 688  
Naphthols in, II. 688, 693  
Nature and origin of, II. 669—70  
Nervous symptoms, II. 704—5  
Oil of eucalyptus in, II. 688  
Perforation of the intestine in, II. 703  
Prophylaxis and disinfection, II. 679—81  
Protective inoculation against, II. 675—76  
Quinine in, II. 688, 690  
Neiapses in, II. 675  
Salicylate of bismuth in, II. 688  
Sailor in, II. 688, 695  
Stimulants in, II. 687—88, 703—4  
Sulphuric acid in, II. 688, 693  
Symptoms of, II. 673  
Thymol in, II. 694  
Treatment of complications of, II. 696  
Treatment of convalescence from, II. 705—6  
Turpentine in, II. 688, 695  
Urotropine in, II. 688, 695  
Vaccine in, II. 676  
See also Additional Formulas, II. 711, 712
- Typhus fever,** II. 707—II  
Course of, II. 707—8  
Diet in, II. 710  
Diffusion of, II. 707  
Hypodermic injections in, II. 710  
Incubation period of, II. 707  
Open-air treatment of, II. 710  
Quinine in, II. 709  
See also Additional Formulas, II. 712
- Ulcer of the Duodenum.** See **STOMACH AND DUODENUM, ULCER OF THE,** I. 71—95
- Uremia,** in Bright's disease, II. 218—19  
General considerations, II. 155, 156
- Urinary and renal affections.**  
See **Albuminuria,** II. 180—85 ;  
**Bright's disease,** II. 205—40 ;  
**Chyluria,** II. 179—80 ; **Cystitis,** II. 200—3 ; **Haematuria,** II. 173—77 ; **Hæmoglobinuria,** II. 177—79 ; **Hydronephrosis,** II. 196—97 ; **Lithiasis,** II. 157—66 ; **Oxainuria,** II. 166—70 ; **Phosphaturia,** II. 170—73 ; **Pyuria and Pyelitis,** II. 197—200 ; **Renal calculi,** II. 187—90 ; **Renal colic,** II. 190—94