

brown shading of three bars on the fascia ; outer margin from the apex to the tails is pure white ; tails are brown, heavily fringed with white. Tails, $\frac{3}{16}$ inch long. Expanse, $2\frac{1}{8}$ inches. Hab.: Bagasas, Costa Rica, Ct. America. Type Ehrmann collection.

Goniurus Cleopatra, n. sp.—Female. Upper side : antennæ, head, thorax and abdomen are dark brown. On the fore wings the ground colour is dark brown ; the costa near the base to the transparent bar that extends across the fascia is white ; the bar that is on the fascia and the spots on the subapical space are the same as in *G. Triptolemus* ; from the median cell the outer margin and tails are white ; fringes white. On the abdominal margin the fringes are scanty and brown. On the hind wings the ground colour is dark brown ; the abdominal margin is dusted with white from the median vein, including the tails. Fringes on the apical and abdominal margin are dark brown.

Under side : palpi are white, dusted with brown ; thorax and abdomen lighter than above ; there is a faint indication of two white lateral bars on the lower side of the abdomen ; legs dark brown. On the fore wings the ground colour is much lighter than the upper side ; the transparent bars and spots are the same ; the inner marginal area is white. On the hind wings the ground colour is very dark brown, slightly dusted with a lighter shade ; outer margin, including the tails, is pure white excepting four faint lunate spots on the apical area. Tails, $\frac{1}{16}$ inch long. Expanse, $1\frac{5}{8}$ inch. Hab.: Suapure, Venezuela. Type in collection of Ehrmann. This species bears considerable resemblance to *G. Orion*, Cram., from the upper Cauca valley of Colombia, S. A., and was represented in the E. A. Klages catch by one fine example, excepting that one hind wing is missing ; otherwise the specimen is perfect and fresh.

Eumesia Pötomoni, n. sp.—Female. Upper side : antennæ black ; head, thorax and abdomen dark velvety-brown. On the fore wings the ground colour is a bronzy-brown. Base and margin slightly darker, and at the end of the di-coidal cell there is a black dot. On the subapical space near the costa there are two minute transparent spots ; then below these dots there is a dark curved line ; this line ends in the lower sub-median cell. The fringes are black. On the hind wings the ground colour is the same as that of the fore wings, but across the fascia are two dark brown bars ; the outer margin is faintly shaded with brown ; abdominal margin light brown ; fringes dark brown.

Under side : palpi and thorax are gray ; abdomen light brown. On both pairs of wings the ground colour is pale brown ; the markings are the same as above. Expanse, $1\frac{3}{4}$ inches. Hab.: Suapure, Venezuela. Type in collection of Ehrmann.