

Also—

R. Aconitæ,	gr. ss	
Veratriæ,	grs. x	
Adipis	3 j.	M.

Sig.—Apply to part.

NEURALGIC HEADACHE.

R. Quiniæ sulph.,	grs. xij.	
Morph. sulph.	gr. j.	M.

Triturate in a mortar, and divide into twelve powders.

Sig.—One powder every four or five hours.

UNCOMPLICATED SUPRA-ORBITAL NEURALGIA.

R. Arsenici,	gr. ¼	
Ext. conii,	gr. j	
Ext. cannabis ind.,	gr. ½	

Sig.—One dose, ter die. *Da Costa.*

OBSTINATE NEURALGIA.

R. Sodæ arseniatis,	gr. ¼	
Cinch. sulph.,	grs. ij	
Conii ext.,	gr. j.	

Sig.—One dose, ter die. During the paroxysm use hypodermic injection of morphia.

CEREBRAL NEURALGIA.

R. Chloral hyd.,	grs. x	
Pot. bromid.,	grs. xx	
Syr. aurant. cort.,	3 ss	
Aquæ,	3 iss	M.

Sig.—One dose at bed time.

Also—

R. Tinct. cinch. comp.,	3 ij	
Fl. ext. cinch.,	j	
Ammon. brom.,	3 ss.	M.

Sig.—One teaspoonful, ter die.

OVARIAN NEURALGIA.

R. Ammon. mur.	3 ij	
Tinct. aconit.,	3 ij	
Syr. aurant. cort.,	3 viij.	M.

Sig.—One drachm, ter die. *Da Costa.*

ANTI-NEURALGIC TONIC.

For long standing cases—

R. Acid. phosphoric, dil.,	gtts. xx	
Tinct. cinch. co.,	3 j	
Strychniæ phos.,	gr. ½	M.

Sig.—One dose, ter die.

VERTIGO (*Gastric*).

R. Argenti oxidi,	gr. ¼	
Capsici pulv.,	gr. ¼	
Ext. colocynth. co.,	gr. j	
Camphor. pulv.,	gr. ½	M.

Ft. pil. No. 1.

Sig.—Take after each meal

IDIOPATHIC VERTIGO.

R. Sodii bromid.,	gr. xv	
Ext. belladonnæ,	gr. ½	
Vin. colch. rad.,	gtt. x	
Muc. acaciæ,	q. s. ft. 3 j.	M.

Sig.—One dose, ter die.

If bowels are costive, take one comp. rhubarb pill each night.

CHRONIC HICCOUGH, FROM NO ASSIGNABLE CAUSE.

R. Tinct. calabar bean,	℥ x	
Pot. carb.,	gr. x	
Mist. acaciæ,	3 j.	M.

Sig.—One dose, ter die.

No vegetables should enter into the diet. The food should be non-stimulant, making use of milk, eggs, etc. In some cases, tinct. of calabar bean alone answers very well. Sometimes a hypodermic injection of morphia cures or relieves.

INCIPIENT SOFTENING OF THE BRAIN,

Attended with bad memory, visions, giddiness and headaches.

R. Acid. phos. dil.,	gtt. xx	
Tinct. cinch. co.,	3 j	
Tinct. nucis vomicæ,	gtt. v.	M.

Sig.—One dose, ter die.

Keep the patient in society; or send him upon a journey. Aim at bringing before his mind new scenes, etc.

SOOTHING NERVINE AND TONIC.

R. Pot. bromid.,	3 ss	
Ferri pyrophos.,	3 ij	
Elix. humuli,		
Aquæ,	aa. 3 iv.	M.

Sig.—Tablespoonful ter die. *McArthur.*

ANODYNE PILLS.

These have the advantage of not affecting the bowels:—

R. Morph. acet.,	grs. ij	
Hyoscyam, ext.,	grs. viij.	M.
Ft. pil. No. xvj.		

Sig.—One pill at bedtime.

LARYNGISMUS STRIDULUS.

Large doses of bromides. For a child two years old, six grains, every two hours, may be given. If there is any source of irritation, as from the stomach or gums, this must be removed. If not complicated with laryngitis, indicated by a hoarse voice, cold sponging is the grand remedy. To arrest a paroxysm, a dash of cold water in the face is often sufficient. If this fail, cold water applied to the whole body is of value. The child should be sponged faithfully and regularly, at least three times a day, and also allow the child to be much in the open air.