

630. *C. caesescens*, L. Wet meadow and marshy places; common.  
var. *vitillis*, Gray. Damp woods; not common.
631. *C. Deweyana*, Schw. Open woods. Bass River, Kent Co.
632. *C. stellulata*, L., var. *scirpoides*, Gray. Wet ground; common.  
var. *angustata*, Gray. Wet ground. Bass River. Rare.
633. *C. scoparia*, Schf. Very common in damp meadows.  
var. *minor*, Boot. Sent from Restigouche by Mr. Chalmers.
634. *C. lagopodioides*, Schk. Moist open woods. Kent Co.
635. *C. adusta*, Boot. Moist borders of woods. Kent Co. There are two forms of this plant as determined by Olney, one of which he identifies as *C. albobutescens*, Schw. var. *glomerata*, Olney, and the other as var. *sparsiflora*, Olney.
636. *C. sœnea*, Willd. var. *sabulonum*, Gray. *C. silicea*, Olney. Sand beaches on the coast.
637. *C. straminea*, Schk. var. *typica*, Gray. Open woods. Kent Co. Var. *tenera*, Gray. Open woods. Kent Co. Rather rare.
638. *C. vulgaris*, Fries. Wet grassy places. Rather rare. A very variable plant and often difficult to determine.
639. *C. aquatilis*, Wahl. Borders of streams or lakes. Common in St. John Co.
640. *C. aperta*, Boot. Wet places; near Richibucto.
641. *C. stricta*, Lam. Wet swampy grounds; Richibucto.
642. *C. lenticularis*, Michx. Wet banks of streams; scarce.
643. *C. salina*, Wahl. Salt marsh at Molus River; only found it once.
644. *C. maritima*, Vahl. Salt marshes; not rare.
645. *C. crinita*, Lam. Banks of brooks; very common.
646. *C. limosa*, L. Peat bogs; common.
647. *C. irrigua*, Smth. Peat bogs; common.
648. *C. Buxbaumii*, Wahl. Boggy grounds; Restigouche.
649. *C. aurea*, Nutt. On limestone rocks; St. John Co.
650. *C. panicea*, L. Wet meadows; Bass River; rare.
651. *C. granularis*, Muhl. Wet grounds; rather rare; Bass River.
652. *C. pallescens*, L., var. *undulata*, Gray. Bass River; rather common.
653. *C. gracillima*, Schw. Damp grounds; Bass River.
654. *C. laxiflora*, Lam. Several forms or varieties of this exceedingly variable plant are found in open woods and copses.
655. *C. umbellata*, Schk. Rocky ground near the Manse at Blackville.
656. *C. Novæ Angliæ*, Schw. St. John. July 1869.
657. *C. Emmonsii*, Dew. Dry bank near Manse, Blackville; June 1865.
658. *C. Pennsylvanica*, Lam. Dry knolls in woods; Kent Co.
659. *C. varia*, Muhl. Dry hills in woods; Bass River.
660. *C. scabrata*, Schw. Wet grounds; Bass River.
661. *C. arctata*, Boot. Margin of dry woods beside Manse, Bass River.
662. *C. debilis*, Michx. Moist grounds; rather common.
663. *C. capillaris*, L. Bass River 1865; rare.
664. *C. flexilis*, Rudge. Moist shady places; Weldford Station.
665. *C. flava* L. Wet meadows; common.
666. *C. Cœderi*, Ehrh. Wet rocks and cliffs; St. John Co.
667. *C. filiformis*, L. Peat bogs and water; Lake Elsie near Richibucto.
668. *C. Houghtonii*, Torr. Dry grass fields; Kent Co.
669. *C. riparia*, Curtis. In water near Weldford Station; 1868.
670. *C. Pseudo-Cyperus*, L. Borders of lakes and in ditches; rather common.
671. *C. hystericina*, Willd. Wet grounds; Lily Lake, St. John.
672. *C. tentaculata*, Muhl. Wet places; common.
673. *C. intumescens*, Rudge. Damp meadows and swamps; common.
674. *C. lupulina*, Muhl. Wet grounds, Hampton.
675. *C. rostrata*, Michx. Bogs. Rothesay.
676. *C. retrorsa*, Schw. Marshy grounds; common.
677. *C. utriculata*, Boot. Wet swamps and water. Kent County.
678. *C. Vaseyi*, Dew. In water at Lake Elsie, near Richibucto.
679. *C. monile*, Tuckerm. Wet places. Bass River. Fredericton.
680. *C. Tuckermani*, Boot. Wet bank of stream, Bass River.
681. *C. oligosperma*, Michx. In a peat bog near Richibucto.
682. *C. miliaris*, Michx. Wet borders of Kennebecasis at Rothesay.

## GRAMINEÆ—Grass Family.

683. *LEERSIA oryzoides*, Swartz. *Rice Cut-grass*. A perennial marsh grass, with widely branched panicles and flat spikelets; the flat leaves and sheaths are covered with minute hooked prickles, which cut the hand when drawn along them; the panicles are also strongly bristly-ciliate. Common in very wet places. Aug.
684. *ALOPECURUS GENICULATUS*, L. *Floating Foxtail Grass*. A perennial grass, bearing a strong resemblance to *Timothy*. The stems are bent at the lower joints; the upper leaves are about the length of the sheaths, and an awn springing from near the base of the panicle, projects about half its length beyond it. Moist grounds. Carleton. July.