

FAMILY COCCINELLIDÆ.

309. *COCCINELLA EPISCOPALIS* Kirby.—Plate v, fig. 4. Length of body 2 lines. Taken in the journey from New York to Cumberland-house.

[229.] Body narrow, nearly linear, having at first sight the aspect of a *Halicta*, underneath black. Head black with three oblong pale yellow spots, two adjoining the eyes on their inner side, and one placed backwards in the vertex; mouth, antennae, and palpi rufous; prothorax and both elytra taken together, pale yellow with two black stripes, common to both, resembling a bishop's crosier, the crook being on the thorax and the stalk on the elytra; suture of the latter black except at the tip; legs pale testaceous; anus, sides of the abdomen, and tips of the ventral segment, except the basal one, pale.

[Belongs to *Nemio* Muls.]

310. *COCCINELLA TREDECIM-PUNCTATA* Linn.—Length of body 3 lines. Several specimens taken in Lat. 54°.

Body oblong, black, lightly and minutely punctured; underneath slightly downy. Mouth and its organs pale rufous; nose white, whiteness with a posterior central lobe; antennae rufous; prothorax white with a large discoidal spot falling short of the anterior margin, where it is truncated; sides lobed, besides which there is a black dot on each side connected with the above spot; elytra reddish-yellow with six largish black dots, namely 1, 2, 2, 1, and one at the scutellum common to both elytra; the first marginal dot is ovate, the rest approaching to round; the tibiae and tarsi are testaceous; there are two transverse white spots on each side the breast, between the four posterior legs; and four triangular pale ones on each side the abdomen.

VARIETY B. Nose rufous, with a parallelogrammiscal white spot between the antennae.

[Belongs to *Hippodamia* Muls.; very common in Canada.]

311. *COCCINELLA TRIDENS* Kirby.—Length of body $2\frac{1}{4}$ lines. Two specimens taken in the Expedition.

[230.] Body rather oblong, very minutely punctured; black underneath, with two distant white spots on the breast, and three contiguous ones on each side of the abdomen. Head black with a transverse white band or trident between the eyes, tricuspidate both anteriorly and