

FAUNE COLEOPTEROLOGIQUE AU MANITOBA

(Continué de la page 175)

HISTERIDÆ

Hister interruptus, Beauv.

" *abbreviatus*, Fab.

" *Lecontei*, Mars.

NITIDULIDÆ

Carpophilus niger, Say.

Colastus truncatus, Rand.

Conotelus obscurus, Er.

Epuræa labilis, Er.

Omosita colon, Linn.

Ips vittatus, Say.

TROGOSITIDÆ

Peltis ferruginea, Lin.

BYRRHIDÆ

Byrrhus americanus, Lec.

HETEROCERIDÆ

Heterocerus mollinus, Kies.

DASYLLIDÆ

Cyphon variabilis, Thunb.

ELATERIDÆ

Adelocera obtecta, Say.

Cryptohypnus bicolor, Esch.

Elater linteus, Say.

" *rubricus*, Say.

" *apicatus*, Say.

Drasterius elegans, Fab.

Agriotes stabilis, Lec.

" *fucosus*, Lec.

" *pubescens*, Melsh.

" *limosus*, Lec.

Dolopius lateralis, Esch.

Melanotus communis, Gyll.

" *fissilis*, Say.

Limoniæ quercinus, Say.

Corymbites resplendens, Esch.

" *spincus*, Lec.

" *merulus*, Lec.