

Sub-division 3. MINORES, Guén.

Family, ERASTRIDÆ, Guén.

Genus 1. ERASTRIA, Ochs.

- 1.
- E. carneola*
- , Guén.—Rare, beginning of August.

Distribution.—United States, (E. Doubleday).

Division 2.—QUADRIFIDEA, Guén.

Tribe, VARIEGATÆ, Guén.

Family, PLUSIDÆ, Boisd.

Genus 1. PLUSIA, Ochs.

- 1.
- P. ærea*
- , Hübn.—July and August.

Distribution.—New York and Florida, (Doubleday); Nova Scotia, (Redman); Orillia, C.W., (Bush).

- 2.
- P. preactionis*
- , Guén.—Common, August.

Distribution.—New York and Philadelphia, (Doubleday); Mass.: (Prof. Shepard); Orillia, C.W., (Bush).

- 3.
- P. mortuorum*
- , Guén.—Common, August.

Distribution.—New York, (Doubleday); Orillia, C.W., (Bush); Sorel and Rouge District.

Tribe, INTRUSÆ, Guén.

Family, AMPHIPYRIDÆ, Guén.

Genus 1, AMPHIPYRA, Ochs.

- 1.
- A. pyramidoides*
- , Guén.—Common in August on the Mountain.

Distribution.—Massachusetts. (Harris).

- 2.
- A. tragogoponis*
- , Linn.—Abundant, beginning of August.

Distribution.—L Original on the Ottawa, (R. Bell). (Eu).

Tribe, EXTENSÆ, Guén.

Family, HOMOPFERIDÆ, Guén.

Genus 1, HOMOPTERA, Boisd.

- 1.
- H. lunata*
- , Drury.—Taken on Treacle in May.

Distribution.—Massachusetts, (Harris); St. Domingo, (B.M.C.)

- 2.
- H. contracta*
- , Walker, M.S.S. N. Sp.—Middle of July.

“*Fœm.* Ferrugineo fusca; palpi arcuati, graciles, ascendentes; alæ lineis nigris undulatis, linea interiore duplicata strigam nigram includente, linea media dentata apud costam dilatata, linea exteriore pallido marginata; anticæ lineis duabus basalibus.”

“*Female.* Ferruginous brown, cinereous beneath. Palpi curved, slender, obliquely ascending, rising higher than the head; third joint lanceolate, about half the length of the second. Wings with black undulating lines; interior line double, the intervening space partly black; middle line dentate, much dilated on the