

in color, with a narrow irregular paler terminal space. Lines lost. Reniform marked in white, constricted, the white scales touching median vein. Hind wings fuscous, with paler fringes and paler at base. No markings beneath. Abdomen pale yellowish gray. Arizona. Coll. B. Neumoegen, Esq.

*Plagiomimicus* Grote.

I called attention some years ago in the Bulletin of the Buffalo Society to the resemblance between this genus and *Polenta*, in the spreading thoracic tuftings, the modification of the clypeus, the shape of the wing and the peculiar pattern of marking, which is so much alike that the two species would be considered congeneric by most observers. I since contradict Mr. Morrison's observation that the fore tibiæ are unarmed in *Polenta*; this was the best distinction. The discovery of new forms confirms me that the genera I have grouped under *Stiriinæ* are all valid, the modifications of structure being apparent and going closely with the pattern. *Stibadium* approaches *Telesilla*, in frosting and sheen (*aureolum*), and leads to this genus as before suggested. With the new species the group (without absolute value) may fall in between *Calpe* and *Plusia*.

*Oxycnemis*, n. g.

Vestiture flattened hairs; a tuft behind thorax of broad curved shining scales widening to their tips; abdomen short, untufted. Size small, body slender. Hadeni-form ornamentation. Eyes naked, unlashd. Fore tibiæ with the whole short broad joint corneous, terminating in a stout central claw or spine; the legs slender, otherwise unarmed. Notwithstanding its Hadeni-form look, the insect appears nearest to *Triocnemis*.

*Oxycnemis advena*, n. s.

♀. Gray, neatly lined, with the bright color and ornamentation recalling *Charadra palata*. T. a. line black, distinct, upright, attached to the large open claviform. Discal stigmata light gray, subequal, with included dots. T. p. line faint. S. t. line black with a following white shade, marked with black on costa, far from margin and easily taken for t. p. line, curved in inferiorly; veins terminally finely marked in black. Hind wings pale fuscous, whitish at base. A white costal shade on primaries above from inception of s. t. line over the disk. Size small. Arizona, Coll. B. Neumoegen, Esq.