

IN view of the growing favor of subcutaneous injections for syphilis, we give the following formula of Sacaze :

R	Metallic mercury.....	20		(.jv.)
	Lanoline	5		(̄jss.)
	Liquid vaseline	35		(̄j. ¼)

Inject subcutaneously from five to seven centigrammes (gr. $\frac{3}{4}$). Repeat every fifteen days.—*Medical Review of Reviews*.

METHYL SALICYLATE IN THE TREATMENT OF GONORRHOEA—By reason of the great power possessed by methyl salicylate of penetrating investing membranes, M. Duquaire (cited in the *Journal des Praticiens*) has conceived the idea that it will reach the gonococci even when they are seated in the deepest layers of the mucous membrane. He reports a case in which its employment cured the disease in five days, but he does not generalize from that one case. He uses the following solution :

R	Methyl salicylate.....	1 part.
	Bismuth subnitrate	20 parts.
	Liquid vaseline	100 parts.

M. These injections are to be given daily. The patient should urinate, then take the injection, and hold it in the urethra as long as possible. The injections are not painful.—*N. Y. Med. Jour.*

FOR TUBERCULAR CYSTITIS.—General treatment should always be instituted, the indications are to improve nutrition, avoiding drugs which upset the appetite. Climate and salt baths may prove of assistance. Banzet advises against large doses of creosote, smaller dosage and persistent exhibition being better. Guyon's favorite formula for giving it is :

R	Creosote	gr. .05.
	Iodoform	gr. .01.
	Sod. arseniat	gr. .001.
	Cynoglosi	gr. .05.

Pulv. benzoin, q. s. to make one pill.

M. Sig. Two pills twice daily at meal times.

General treatment may in some few instances prove even curative. Instances are cited from the practice of Guyon which go to show in a positive manner the efficacy of general treatment. Banzet warns against putting such cases on a low diet because of a slight albuminuria, a treatment capable of rendering this class of patients much worse.—*Journal Cutaneous and Genito-Urinary Diseases*.