

GROUP WITH MODERATE PALPI.

- | | | |
|--|---|--|
| 8. <i>Hyd. autumnalis</i> Ström
or <i>pluviata</i> Gn. | { | White mesial space, olive shadings,
slate-coloured bands. |
| Var. (a) <i>perfracta</i> Swett..... | { | (a) White mesial space suffused
with pink. |
| Var. (b) <i>crokeri</i> Swett..... | { | (b) Green ground colour replaced
with yellow, black bands. |
| Var. (c) <i>columbiata</i> Taylor..... | { | (c) As typical, mesial space suffused
with cinereous and narrow, very
large size. |
| Var. (d) <i>constricta</i> Strand..... | { | (d) Mesial lines smutty, lines in-
distinct. |
| Var. (e) <i>nigrescens</i> Hoyn.-Hewne.. | { | (e) Unicolorous; smutty suffused. |
| 9. <i>Hyd. transfigurata</i> Swett..... | { | Green ground colour, tendency of
extra- and intradiscal lines to join
near inner margin. |
| 10. <i>Hyd. californiata</i> Pack..... | { | Red shadings to lines and mesial
space. |
| 11. <i>Hyd. lanavahrata</i> Streck..... | { | Said to be var. of <i>californiata</i> . |
| 12. <i>Hyd. glenwoodata</i> Swett.... | { | Long-winged slender species with
s-shaped intradiscal line. |
| 13. <i>Hyd. magnificata</i> Taylor..... | { | Black and green. |

I wish to thank Mr. A. F. Winn and Mr. G. Chagnon for the loan of specimens, and the latter for his help on the genitalia, on which we hope to publish something later. I append a few of the more important references to *Hyd. autumnalis* and its varieties:

1783. *autumnalis* Ström, Kgl. dansk. Vid. Selsk. Skr., p. 85.
 1776. " Dennis & Schiff, Syst. Verz. Wien, 109, 5 (not described).
 1786. *autumnalis* Sepp., II, pl. 5, figs. 1-8.
 1794. *trifasciata* Bork., Eur. Schmett., V, p. 308, No. 141.
 1797. *impluviata* Hüb., Eur. Schmett., 223 (post 1797).
 1810. " Haw., Lepid. Brit, part 2, p. 321.
 1822. " Hüb., Verz. Schmett., 322, 3, p. 106.
 1828. " Treits., Schmett. Europa, VI, part 2, p. 21.
 1829. " Stephens, Nomenc. Brit. Ins., p. 44.
 1830. " Dupon., Lep. France, V, p. 424, pl. 200, fig. 3.
 1831. " Stephens, Ill. Brit. Haust., III, p. 254.