

minently raised white veins and a thin indument of short white hairs; panicle of fertile plant small but rather many-flowered and dense, its branches glabrous, as also the calyx; pistils numerous, encircled by a row of 3 to 7 or 8 stamens; filaments abruptly clavate above, but not to the width of the mostly broad and merely oval (occasionally oblong) anthers; immature carpels fusiform, substipitate, glabrous.

The type specimen of this rather elegant and very pale meadow-rue is in the herbarium of the Geol. Survey, under No. 869, and came from Tignish, at the northern extremity of Prince Edward Island, where it was obtained by Professor Macoun, 25 July, 1888. The flowers, though mainly pistillate, are very fairly hermaphrodite.

A plant much like this in habit, quite as pale with bloom, and even with very similar foliage, is in the same herbarium from Boyleston, Nova Scotia, by Dr. Charles A. Hamilton, July, 1890. It also has hermaphrodite flowers, and in these the calyx is persistent and is of five or six narrow elliptical sepals. This plant also I refer tentatively to *T. glaucodeum*, and hope it may prove to belong with the more northerly and insular type.

BIRDS OBSERVED AT SHERBROOKE, QUE.—SPRING MIGRATION, 1909.

BY R. G. PRICE.

Crow.....	Feb. 28, common.....	S.R.
Robin.....	Mar. 12, ".....	S.R.
Horned Lark.....	" 12, fairly common (1)....	S.R.?
Bronze Grackle.....	Apr. 5, common.....	S.R.
Bluebird.....	" 5, ".....	S.R.
Red-tailed Hawk.....	" 5, becoming rare.....	S.R.
Red-shouldered Hawk....	" 6, common.....	S.R.
Song Sparrow.....	" 6, ".....	S.R.
Marsh Hawk.....	" 5, fairly common.....	S.R.
Northern Shrike.....	" 6, ".....	R.
Blue Jay.....	" 6, common.....	R.
Red Crossbill.....	" 6, fairly common (2)....	P.M.
White-winged Crossbill...	" 6, rare.....	P.M.
Pigeon Hawk.....	" 7, fairly common.....	S.R.
Junco.....	" 7, ".....	P.M.
Meadow Lark.....	" 12, ".....	S.R.
Sparrow Hawk.....	" 12, ".....	S.R.
Golden-eye Duck.....	" 12, common (3).....	S.R.
Cowbird.....	" 15, ".....	S.R.