

the "Ottawa Cow." Where trees have become diseased from any cause, or where a fire has ravaged a pine forest and scorched and partially destroyed the timber, or where logs after being cut have been allowed to remain a season in the woods or in the mill yard—there these insects gather and soon multiply to a prodigious extent. The mature insect is over an inch in length; the antennæ of the male reaches the extraordinary length of from two to three inches, while those of the female are shorter. The female lays her eggs in the crevices of the bark, where the larvæ when hatched eat their way into the wood, burrowing extensive galleries through the solid timber; when mature they are large, white, almost cylindrical, footless grubs. They pass their chrysalis stage within their burrows, and the perfect insect on its escape eats its way out through the bark. There are about a dozen species in this family known to be destructive to pine.

Most of the insects belonging to the family *Buprestidæ* may be recognized by their brilliant metallic colors¹; they have very short antennæ which are notched on one side like the teeth of a saw, and are often hidden from view by being bent under the thorax. *Chalcophora liberta* is one of the most destructive to pine trees, and its history is very similar to that of the long-horned beetle just described, but the larva is of a different form, and has the anterior segments or rings of the body very large, reminding one of the appearance of a tadpole. The perfect insect is about three-quarters of an inch long, of a brassy or coppery hue, with the thorax and wing-covers deeply furrowed by irregular longitudinal depressions. Dr. Fitch enumerates twelve species belonging to this family which are known to be injurious to pine. Additional information in reference to these beetles may be found in an article contained in the last annual report of our Society, by Mr. J. Fletcher, of Ottawa.

The cylindrical bark beetles, *Scolytidæ*, are also a numerous family, eight species of which are known to attack pine. The boring *Hylurgus*, *Hylurgus terebrans*, is probably one of the commonest. This beetle is about a quarter of an inch long, of a chestnut red color, thinly clothed with yellowish hairs, and is found during the month of May. The larva, which is a small yellowish white footless grub, bores winding passages in every direction in the inner layers of the bark of the tree, and also through the outer surface of the wood.

In some parts of our Province pines are greatly injured and sometimes killed by the attacks of a woolly bark louse, which covers parts of the