

BAYER'S MAGEUTICAL PRODUCTS

A tasteless, odour-less, nutrient meat powder; it contains all the albuminoid principles of the meat in an easily soluble form. It has been extensively employed and found to be of the greatest service in consumption, diseases of the stomach and intestinal tract, chlorosis and rick-ets. It is of great value in convales-cence from all diseases. SOMATOSE strengthens the muscles and stimulates the appetite in a remarkable manner. Dose for adults: a level teaspoonful three to four times a day with milk, gruel, coffee, etc.

(Ferro-IRON SOMATOSE Somatose).

A first-class tonic, containing the albuminous substances of the nieat (albumoses) organically combined with iron. Special indications: Chlorosis and Anaemia. Daily dose: 75 to 150 grains.

MILK-SOMATOSE (Lacto-Somatose).

A strength giving food containing the albuminous matter (albumoses) of the Solutions of 1/4 to 2%. Ointments. milk.

TRIONAL A most reliable and quickly-acting hypnotic of the Sulfonal group. Dose: 16 to 20 grains, in a large cup of hot liquid.

The active prin-IODOTHYRINE ciple of the thyroid gland. It is most efficacious in Strumous Diseases, Myxoedema, Obesity, Rickets, Psoriasis, Eczema, and Uterine Haemorrhages. Dose: grains two to eight times a day for adults; 5 grains one to three times daily for children.

LYCETOL

PHENACETINE-BAYER PIPERAZINE-BAYER

Solvent. Has a marked effect on the diuresis. Dose: 16 to 32 grains daily.

An Iodine Cicatrisant ARISTOL which is an excellent odourless substitute for Iodoform and highly recommended for Burns, Wounds, Scrofulous Ulcerations, etc.

EUROPHEN A perfect substitute for Iodoform. Odourless and non-toxic. Has a covering power five times greater than Iodo-form. Especially useful in Ulcus molle et durum.

cresoltriiodide OSOPHAN. particularly efficacious in the treatment of all kinds of cutaneous disorders caused by animal parasites.

A new silver pre-PROTARGOL paration. Most reliable in cases of Gonorrhoea. Antiseptic wound healer. Excellent results in cases of Gonorrhoeal Ophtalmia.

TANNIGEN. An almost tasteless intestinal astringent. Most efficacious in Chronic, Acute and Summer Diarrhoeas. Adult dose: 8 grains every three hours.

TANNOPINE (Formerly none"). new Special Indicaintestinal astringent. tions: Tuberculous and non-tuberculous Enteritis, Typhus. Dose: 15 grains, three or four times daily.

SALOPHEN Specific for Influenza, Headache, Migraine, Acuté Articular Rheumatism, Chorca, SALOPHEN Tartrate of Piperazine Sciatica. Dose: 15 grains four to six Anti-Arthritic, Uric times daily. In powders, etc.

SULFONAL-BAYER SALOL-BAYER

ples and literature may be had on application to the

DOMINION DYEWOOD & CHEMICAL CO., TORONTO.

Sole Agency and Depot in Canada for all " BAYER'S" Planniquestical Products.

(WHOLESALE ONLY)