

21. *T. PURPURASCENS* (Purplish Meadow-rue).

Templeton and along the Ottawa below the city (H. M. Ami*),
Hull (J. F.) This species is distinguishable from *T. Cornuti*
by having the anthers drooping on capillary filaments and the
flowers greenish purple. On the other hand, *T. Cornuti* has
white flowers, short filaments and anthers not drooping.

22. *T. CORNUTI*, L. (White-flowered Meadow-rue).

Islands and wet meadows and swamps; a tall, handsome plant.
July—1. (B.)

RANUNCULUS, L. Crow-foot. Buttercup.28. *R. AQUATILIS*, L. *v.* *TRICHOPHYLLUS*, Chaix (White Water Crow-foot).

Stagnant and slow-flowing waters. Ju.—1.

29. *R. MULTIFIDUS*, Pursh (Yellow Water-Crow-foot).

In shallow water, and creeping over mud, Malloch's Bay, Dow's
Swamp and Kettle Island (J. F.), Billings Bridge (R. B. Whyte).
Ju.—2.

Var. *y. REPENS*, Watson.

Leaves all round-reniform, palmately, 3—8 cleft; stems creeping
in mud, Casselman and Gatineau Point (Prof. Macconn). Ju.—2.

31. *R. FLAMMULA*, L. *var.* *REPTANS*, Meyer (Smaller Creeping Crow-foot).

Mud-flats and river margins July—1.

32. *R. CYMBALARIA*, Pursh (Salt-water Crow-foot).

In a low, wet meadow at Thurso, P.Q. Ju.—2.

37. *R. ABORTIVUS*, L. *var.* *MICRANTHUS*, Gray (Small-flowered Buttercup).

Woods and Meadows. From "Macconn's Catalogue," Part III.,
p. 480, it would appear that our Ottawa plant is not the type of
R. abortivus, L. May—2. (B.)

30. *R. RECURVATUS*, Poir (Downy Wood-Buttercup).

Woods. An attractive species with large leaves and small flowers.
May—2. (B.)

45. *R. acris*, L. (Meadow Buttercup).

Introduced from Europe. Very abundant in meadows. June (B).