

but such artificial detachment is usually incomplete, is liable to injure the uterine tissue, and the operator's hand may be the bearer of septic germs, or these may pass in with the air admitted during the manipulation, and find a congenial soil for their development in fragments of placenta, or blood-clots that are retained in the uterus. Therefore, unless hemorrhage demands immediate interference, the obstetrician refrains from passing his hand into the uterine cavity for the removal of an attached placenta; a completely adherent placenta is not so dangerous as the intra-uterine use of the hand for its detachment. I believe, then, that armed expectation is wise in the latter case, only endeavoring, by suitable compression of the uterus with the hand actually through the abdominal wall, to determine or assist that retraction of the organ which is nature's method of separating the placenta. After the detachment of the placenta—a fact which is best learned by feeling part of the organ with the finger passed into the mouth of the womb—we may, by friction and compression of the uterus, if needed, evoke uterine contractions which will cause its expulsion. Those who believe that the placenta presents its foetal surface at the os uteri, urge the value of moderate and continuous traction upon the cord, thus assisting the moulding of the mass to the orifice through which it is to come. This conservative view as to the management of so-called retained placenta has been strongly presented by Siredey in his recent work upon puerperal diseases. The common expression, retention of the placenta, means very different conditions, each requiring its appropriate treatment.

Dr. Parvin's report contains another important lesson on septicæmia and its temperature and the difficulty of diagnosing a case of septicæmia from malarial infection, and concludes with a brief study of a ruptured uterus.

The uterus was ruptured in consequence of a shoulder presentation, a case which ended in death the eighth day after delivery. Yet I would fail in duty to my profession that has been so good, so generous to me, if I did not make the case fully known. The patient was a well-formed healthy multipara; she had been in labor nearly twelve hours when I first saw her, the left shoulder presenting. Ether was immediately given until she was thoroughly under its anæsthetic effect; and then, without violence, nay, with great ease, I passed two fingers behind the right knee, brought the foot down, and turning and delivery were effected in a few minutes; the placenta followed almost immediately; the child, quite a large one, was dead. The patient came out from the anæsthesia satisfactorily; her pulse was good; there was no complaint, no shock, no great hemorrhage. Yet that woman had a ruptured womb, the tear beginning at the os uteri on the right side, involving the cervix and the lower part of the body of the uterus, this condition being made known by the post-mortem. If it be thought I ought to have

known this accident at the time of delivery, I can only say that like ignorance happened to Dubois, to Hervieux, to Tarnier, and others—the first revelation of the uterine rent being made at the post-mortem; these silent tears of the womb are, as Hervieux has suggested, probably more frequent than generally thought. No, my self-reproach is not in this, but in not having made myself, or by another, an examination during pregnancy, so that the abnormal presentation could have been corrected, if not then, at least early in labor. But let this pass. The great practical lesson to be drawn from the accident is not only the importance of an early rectification of a malpresentation, but also an appreciation of the danger of rupture of the uterus and how this accident occurs. The drawing now shown gives the position occupied by the child, and also and especially gives the change in form and thickness of the two cavities of the uterus, which, as so admirably described by Bandl, are formed when nature is unable to overcome the obstacle to labor found in such case. The one cavity is formed by the body of the uterus, and its walls become thicker and stronger; the other, by the cervix, and its walls grow thinner—become indeed so attenuated and weak that a very slight additional strain at some point; that strain may come from a uterine contraction, or solely from the introduction of the finger; and thus peril from action, peril from delay must be before the obstetrician's mind when called to a case of neglected shoulder presentation.

Of course had I seen this patient an hour or two earlier, the event might have been different. The pressure of the presenting part had been so severe that a slough of the vesico-vaginal wall occurred, and the patient, had she had recovered, would have required an operation for the resulting urinary fistula; I have thought that possibly the uterine rent was in part the result of a slough also; but be this as it may, there was not the slightest indication given at the post-mortem that any hemorrhage in the abdominal cavity had taken place.—*Chicago Medical Press.*

TREATMENT OF ECZEMA OF THE GENITALIA; AND LEUCORRHEA.

In cases of eczema, in which glyceroles and unguents have failed, the following formula has been successful:

Chlorate of potassium.....30 grains;
Wine of opium.....50 grains;
Pure water.....1 quart.

Applied to the parts by linen compresses covered with oiled silk. If there is much inflammation, precede this with warm hip-baths and cataplasms sprinkled with powdered carbonate of lime. In obstinate pruritus, associated with leucorrhœa, a tablespoonful of a mixture of equal parts of tincture of iodine and iodide of potassium; in a quart of warm tar-water (tar-water holding the iodine in