

A good local anesthetic for spraying abscesses before lancing is made with half a drachm of chloroform to $\frac{3}{4}$ of ether.

Turpentine is recommended as a deoderant for the hands and instruments after iodoform has been used.

A tablespoonful of cream before meals is recommended for congenital constipation.

Ten drops of a 6 per cent. solution of cocaine injected into a carbuncle at the incipient stage will often abort it.

When albumen is present during pregnancy, put the patient on milk diet, see that the bowels are kept open and give a warm bath every three or four days.

Examine the urine for sugar in all cases of carbuncle and eczema—especially eczema of the genitals.

A Burgundy Pitch plaster is often of service to females who complain of weakness and pain in the lumbar region. It stimulates the surface, excites a pleasant glow and gives mechanical support.

Nitrate of silver stains can be removed from the skin by painting them with iodine, and then applying liq. ammonia B. P. or even spts. ammon. aromat.

Inunctions of cod liver oil, or even olive oil, are beneficial in the wasting diseases of children.

It is said that the beneficial result in goitre from the use of the thyroid extract is due to the iodine which is in it, and that the disease results from a deficiency of iodine in the system, and that the drug alone is as good as the extract.

A little water may be safely administered to an infant to quench thirst, as mother's milk does not do it. Infant often cry simply from thirst.

In collapse a hypodermic injection of a $\frac{1}{4}$ gr. of Morph. with the $\frac{1}{100}$ of a grain of Atropine is an excellent stimulant.

Spirits of camphor applied to a carbolic acid burn will arrest the pain in a few seconds.

Pepto-Mangan (Gude) is used largely by Dr. F. W. Campbell in cases of Anemia and he says it seldom fails to give very excellent results.