

E, tightly over the shield, and straightening the brace from time to time, the deformity is gradually and gently reduced. At each re-application the brace is made a little straighter than the foot at that stage. This may readily be done by the hands, and then the adhesive strip is to be tightened over the shield, till the shape of the foot agrees with that of the brace. After a few days,




Fig. 5.

Fig. 6.

the brace is to be made still straighter, and again re-applied, and made tight till another point of improvement is gained. The brace is applied very crooked at the beginning of treatment, as in Figs. 3 and 4, and is straightened from time to time, and a longer brace applied as the deformity is reduced and the patient grows. It should be removed every week, or two weeks, and an interval of a few days allowed for freedom from the brace, when the mother is advised to manipulate the foot constantly, using as much force as she will in the direction of symmetry. Manipulating the foot during these intervals is of great importance, as cases have occurred in which varus and equinus have been entirely overcome by the mother's hand alone.

By this simple and prosy treatment, carried out systematically and without haste, or violence, or pain, the foot, unless it is a frightful exception, may, with certainty, be changed from varus to valgus. At the same time, the tendo-achillis is lengthened till the position of the foot is near the norme, or at right angles with the leg, as the result of manipulation and giving the brace from time to time




Fig. 7.

Fig. 8.

a partly antero-posterior action. Figs. 3 and 4 show approximately the shape of the brace at the beginning of treatment, Figs. 5 and 6 when the varus is reduced, and Figs. 7 and 8 when valgus has taken the place of varus. The foot, in this latter stage, may not hold itself valgus, when left to itself, but, with almost no force, and with one finger, it may be pushed into valgus; and in this condition it must

be when the child begins to walk, and then another stage of treatment begins.

When the patient begins to walk we have a new difficulty. It is now seen that the weight of the body, falling on the tender and ill-formed foot, will, if not properly directed, defeat all our efforts. Let us, for a moment, consider the mechanical environment of the human foot. In the first place, the corporal weight, which the quadruped distributes among four pedal extremities, falls, in man, upon two. Again, the small floor area covered by the feet and their slight structure, seem unequal to the task of supporting the towering frame above them, which in some cases almost resembles a pyramid resting on its apex. And when we observe the effect of active locomotion we see weight and momentum combine in an apparent effort to crush and destroy. And, furthermore, when extraneous weights are added and the strain is prolonged, as in the case of the burden-bearer among savage tribes, or the infantry soldier on a forced march, the endurance of the foot excites wonder. It is not strange that the feet are subject to ailments: to blisters, bunions, ingrowing nails, hallux valgus, hammer toes, loss of the arch, weak ankles, painful affections of the metatarsus, perforating ulcers, osteitis, and the varieties of talipes. The wonder is that they are not permanently disabled soon after walking is begun, and certainly when the adipose tissue of the body takes on the development which accompanies age and good living. The gourmand, Savarin, said that, among the works of creation, the design of the human foot was a conspicuous failure. Considering the immense weight carried by the foot, it is evident, however, that only the most perfect natural adaptation of mechanics has enabled this insignificant member to perform its superlative functions, and that great caution should attend all procedures having for their object its artificial re-construction.

It is also sufficiently evident that the correction of club-foot by mechanical means, while the patient continues walking, is a problem beset with difficulty. We have, however, a luminous ray of hope and encouragement in the observation that, in talipes varus, there is an important boundary line between deformity and the norme. If the foot is held in some way, now to be considered, on the right side of this boundary line, each step forces