- 6. Lestes disjuncta, Selys.—Toronto, July 15-Aug., 1904, abundant along the banks of the Humber River and in neighbouring woods; Toronto Island, July 25, 1904, flying over the lagoons; Lake Simcoe, Aug., abundant along the margins of creeks, Algonquin Park, Aug. 13, 1903.
- 7. Lestes rectangularis, Say.—Chatham, Aug. 10, 1901; Niagara Glen, June 28, 1903; Toronto, June 24, 1904, Aug. 9, 1903; De Grassi Pt., July-Aug.; Algonquin Park, Aug. 11-29, 1903. A common but not abundant species.
- 8. Lestes vigilax.—Sarnia, Aug. 8, 1901; Grenadier Pond, Toronto, late June-Aug. A very local species, exceedingly abundant around Grenadier Pond. I have reared the flies from nymphs taken from the margins of the pond.

Sub-family Agrionina.

- 9. Argia putrida, Hagen.—Algonquin Park, July 15, 1900, one teneral (J. Macoun), Aug. 15-21, 1904 (P. Hahn). I found the exuviae on a timber slide at Ragged Lake, Algonquin Park, Aug., 1902. This species will doubtless be found commonly along the shore of Lake Erie.
- 10. Argia apicalis, Say.—Chatham, Aug. 10, 1901; abundant along the banks of the Thames.
 - 11. Argia violacea, Hag. Algonquin Park, Aug. 17, 1903.
- 12. Chromagrion conditum, Charp Berlin, Aug., numerous (W. J. Fraser).
- 13. Nehalennia irene, Hag. Toronto, June 22-July, common, especially around Grenadier Pond; Rosebank, June, 1903; De Grassi Pt., July-Aug. 3, 1903.
- 14. Amphiagrion saucium, Burm.—Reported from Ontario by Calvert (Cat. Odon. Phil., 236, 1803).
- 15. Enallagma Hageni, Walsh.—Toronto, June 9-July; Rosebank, June; De Grassi Pt., July; Go Home, Georgian Bay, July 18, 1904; Algonquin Park, Aug. 17-28, 1904. Our commonest Enallagma, occurring in swarms around Grenadier and other ponds in June.
- Enallagma geminatum, Kellicott.—Toronto, Grenadier Pond, July 11-14, 1904. Not common.
- 17. Enallagma exsulans, Hag. Chatham, Aug. 10, 1901; De Grassi Pt., July 18, 1904; Algonquin Park, Aug. 24, 1002.
- 18. Enallagma ebrium, Hag.—Toronto, June 3-14, 1901, Aug., abundant; De Grassi Pt., July 3, 1904, common; Algonquin Park, Aug., 1904 (P. Hahn). Nearly as common as E. Hageni, except northward.