







Concentrations of carbon dioxide and methane after remaining relatively constant up to the 18th century, have risen sharply since then due to man's activities. Concentrations of nitrous oxide have increased since the mid-18th century, especially in the last few decades. CFCs were not present in the atmosphere before the 1930s.

The effect of ozone on climate is strongest in the upper troposphere and lower stratosphere. Model calculations indicate that ozone in the upper troposphere should have increased due to human-made emissions of nitrogen oxides, hydrocarbons and carbon monoxide. While at ground level ozone has increased in the northern hemisphere in response to these emissions, observations are insufficient to confirm the expected increase in the upper troposphere. The lack of adequate observations prevents us from accurately quantifying the climatic effect of changes in tropospheric ozone.

In the lower stratosphere at high southern latitudes ozone has decreased considerably due to the effects of CFCs, and there are indications of a global-scale decrease which, while not understood, may also be due to CFCs. These observed decreases should act to cool the earth's surface, thus providing a small offset to the predicted warming produced by the other

greenhouse gases. Further reductions in lower stratospheric ozone are possible during the next few decades as the atmospheric abundances of CFCs continue to increase.

Concentrations, lifetimes and stabilisation of the gases

In order to calculate the atmospheric concentrations of carbon dioxide which will result from human-made emissions we use computer models which incorporate details of the emissions and which include representations of the transfer of carbon dioxide between the atmosphere, oceans and terrestrial biosphere. For the other greenhouse gases, models which incorporate the effects of chemical reactions in the atmosphere are employed.