

phenylpropanolamine
 pipradol
 prolintane
 propylhexedrine
 pyrovalerone
 strychnine

and related compounds

B. Narcotic analgesics e.g.:

alphaprodine
 anileridine
 buprenorphine
 codeine
 dextromoramide
 dextropropoxyphene
 diamorphine (heroin)
 dihydrocodeine
 dipipanone
 ethoheptazine
 ethylmorphine
 levorphanol
 methadone
 morphine
 nalbuphine
 pentazocine
 pethidine
 phenazocine
 trimeperidine

and related compounds

C. Anabolic steroids e.g.:

bolasterone
 boldenone
 clostebol
 dehydrochlormethyltestosterone
 fluoxymesterone
 mesterolone
 metandienone
 metenolone
 methyltestosterone
 nandrolone
 norethandrolone
 oxandrolone
 oxymesterone
 oxymetholone
 stanozolol
 testosterone*

and related compounds

D. Beta-blockers e.g.:

acebutolol
 alprenolol
 atenolol
 labetalol
 metoprolol
 nadolol
 oxprenolol
 propranolol
 sotalol

and related compounds

* For testosterone, the definition of a positive depends upon the following: — the administration of testosterone or the use of any other manipulation having the result of increasing the ratio in urine of testosterone/epi-testosterone to above 6.