described effect is attributable to the high absorbability of this preparation as compared with the numerous other chalybeates, and, further, to the combined action of iron and manganese upon the blood-forming organs. I would add that numerous investigators, such as Hannan, Kugler, and many other authors, have called attention to the important part played by manganese both in the blood and as a hematogenic remedy.

In the case under consideration there was a perceptible improvement in the patient's subjective and objective state. The existing disturbances subsided gradually; the cardiac palpitation, loss of appetite, and sleeplessness disappeared, and after four weeks' treatment she was discharged cured.

It is not the purpose of this report to detail numerous histories of cases, and I shall content myself with briefly mentioning that I have treated more than 100 cases of chlorosis with Gude's Pepto-Mangan with as good results as those above described, except that in some instances the results did not appear as promptly. The fact cannot be sufficiently emphasized that during the entire course of treatment the remedy did not have to be discontinued on a single occasion, although this must be often done with other ferruginous preparations. I never heard a complaint that the preparation was not well tolerated; on the contrary, the patients stated that they did not experience the slightest disturbance even during its prolonged use, and that it acted mildly, was well borne, caused no disturbance of digestion, but rather promoted the latter, and was free from any disagreeable taste.

I have previously mentioned that it may be positively assumed that Pepto-Mangan stimulates the hematopoietic organs to increased activity. Numerous blood findings discovered casually by me, the appearance of the so-called immature forms of blood corpuscles, constrain me to take this view. Of much greater importance is the circumstance, however, that in numerous diseases of the blood occurring in connection with the lymphatic and blood-making organs, I have derived excellent results from Gude's Pepto-Mangan.

Decided amelioration in the leukemic state, arrest of the process in severe cases for a long time, reduction of the glandular swellings, improvement in the relation between the red and white corpuscles, were noted by me in several cases under my care.

In my opinion, the value of ferruginous preparations in neurasthenia and hysteria has received too little consideration. The success of a rational therapy depends upon an effective application of all methods of treatment and remedies which enable us to combat the entire group of symptoms. An easily absorbable