HEADACHE POWDERS.

Directions for use as found on the label.	Remark		RESULTS OF ANALYLIS.					
		rks.	Weight of Con- tents of one pow- der, grain.	Acetanilide = Anti-febrin, grains.	Phenacetin, grains.	Phenazone = Anti- pyrine, grains.	No. of Sample.	Remarks and Opinion of the Chief Analyst.
G. FERGUSON, IN	SPECTOR	-Conclude	d.					
taken. A seidlitz pow- der or other mild lax- ative should be taken with the first dose. Care should be taken in administering to children. Reduce dose according to age. Dose for sleeplessness, one powder on retiring. These powders may be taken dry on the taken dry on the								
K. X. W. E. BELAND	, INSPEC	TOR.						
birections: One powder every three hours; however, the dose can be repeated after one hour if not relieved. For children 1 to 2 years old, divide one powder in 12; 3 to 4 years, divide one pow- der in 8; 5 to 10 years, divide one powder in 4; 10 to 15 years, divide one powder in 2. How to take them: Place a powder on the tongue and drink a little water		0. 1	6.8	4.9	none.	none.	36391	Curative propertie claimed. No declara- tion of acetanilde- violates requirement of P. or P. Ac- acetanilde being in excessive amount.
after, privections:Placea pow- der on the tongue and swallow with a draught of water. If not re- lieved in half an hour take another powder.		p. 1 2	11-5	4.9	11	"	36392	Curative powers claim ed. No declaration of acetanilide—violate P. or P. Act, a acetanilide in exce- sive amount; but i protected by stamp
	Powder No	1	8.3	1.7		None.		Curative powers clain ed. Meets require
***************	Powder No		5.0	2:0 2:5	None.	None.	36394	ments of P. or P. Act Acetanilide Declared Meets requirement of P. or P. Act