

ham, Proc. Zool. Soc., London, 1897, 139-175). The following synopsis of genera is from Walsingham, with addition of two genera described since the original publication :

ANAPHORINÆ.

Palpi erect or slightly recurved.

Veins 8 and 9 of fore wing stalked.

Palpi erect.....*Eulepiste*.

Palpi appressed to head.....*Neolophus*.

Veins 8 and 9 separate.

Antennæ bipectinate.....*Ankistrophorus*.

Antennæ simple or serrate towards apex.

Tarsal joints of hind legs strongly fringed above...*Thysanoskelis*.

Tarsal joints not fringed above.

Palpi erect, with separate tufts on each joint.....*Ortholophus*.

Palpi slightly recurved, uniformly hirsute.....*Pseudanaphora*.

Palpi strongly recurved.

Antennæ bipectinate.....*Felderia*.

Antennæ serrate throughout.

Veins 8 and 9 of fore wings stalked.....*Cænogenes*.

Veins 8 and 9 of fore wings separate.....*Anaphora*.

Antennæ simple, compressed, or slightly serrate at ends.

Veins 7 and 8 of fore wings stalked.....*Atopocera*.

Veins 8 and 9 of fore wings stalked.

Head with an erect crest.....*Urbara*.

Head without an erect crest.....*Hypoclopus*.

No veins of fore wings stalked.

Palpi roughly clothed throughout.

An erect fringe along lower margin of cell on hind

wings.....*Pilanaphora*.

No such erect fringe on hind wings.....*Acrolophus*.

Palpi smooth, the last joint only tufted.....*Stæberhinus*.

Genus EULEPISTE, Walsingham.

Wals., Trans. Am. Ent. Soc., X., 169, 1882; Trans. Ent. Soc., Lond., 1887, 142.

Synopsis of Species.

Uncus single, the opposing lower limb not half as long as the upper limb.

Harpes concave, rounded at the ends.....*Cressoni*.

Harpes concave, obliquely truncate above.....*maculifer*.