

the second segment and the apex of the first bright yellow; flagellar segments with the basal enlargement very prominent, blackened, the remainder of the segments dark brown, the apical segments more uniform; antennæ elongated for this group of species. Head brownish gray, clearer and brighter toward the occiput and around the eyes; the vertex with a low tubercle bearing an impressed median line; a circular to transverse impressed mark just behind the base of the antennæ meeting on the middle line before the tubercle.

Mesonotal præscutum light gray, the stripes darker gray to brown, narrowly margined with dark brown; the median stripe is bisected by a pale line, which is likewise margined with brown; scutum brown with two light gray spots on each lobe, the largest lying caudad and proximad; scutellum and postnotum light gray, with a narrow brown median line. Pleura with the dorso-pleural membrane dull yellow, the sclerites light gray with a rounded brown spot just beneath the anterior spiracle and a smaller one on the mesopleura. Halteres light yellow, the knob brown. Legs with the coxæ dusted with light gray; trochanters brownish yellow; femora brown, the apex darker; tibiæ brown, lightest at the base; tarsi brown. Wings with a light gray tinge, the costal cell more yellowish, stigma brown; a brownish blotch at the arculus and the origin of *Rs*; veins and the deflections of veins narrowly seamed with brown; hyaline spots before the cord, beyond the stigma in cell *2nd R₁*, base of *R₂* and *R₃*; a rounded blotch in cell *M* at two-thirds the length of the cell; pale flecks in cell *1st A*.

Abdominal tergites varying from almost clear yellow throughout to a brownish yellow, the caudal margin narrowly yellowish, the lateral margins narrowly grayish; the base of the tergites with a transverse rectangular coarsely punctured area on either side of segments 3 to 5, somewhat as in *T. discolor* Loew and similar species; second sternite brownish yellow with a large rounded black spot at the base; segments two to five with punctured areas on the sides of the sclerites, on segments two, four and five transverse, on segment three oblique; apical sclerites brown, broadly yellowish on the caudal margin. Ninth tergite yellowish, the caudal margin with a small, shallow, rounded or rectangular median notch, the lateral lobes broad, obliquely truncated; dorsal surface with a