We should be glad to have you write for a sample of

TAKA-DIASTASE.

Acts more vigorously on starch than does Pepsin on proteids.

RELIEVES Starch Dyspepsia.

We are now able to relieve a large number of persons suffering from faulty digestion of starch, and can aid our patients, during convalescence, so that they speedily regain their weight and strength by the ingestion of large quantities of the heretofore indigestible, but nevertheless very necessary, starchy foods. We trust that the readers of the Gazette will at once give this interesting ferment a thorough trial, administering it in the dose of from 1 to 5 grains, which is best given in powder, or, if the patient objects to the powder, in capsule — The Therapeutic Gazette.

Pepsin is of No Value

In ailments arising from

Faulty Digestion of Starch.

PARKE, DAVIS & CO.,

ずたみたみたみたみたみたみたみたみたみたみたんたたんだた

BRANCHES :

~~ 必然がなればればればればればればればれば

NANCHES.

NEW YORK: 90 Maiden Lane.

KANSASCITY: 1008 Broadway.

BALTIMORE: 8 South Howard St.

NEW ORLEANS: Tchoupitoulas and Gravier Sts.

BRANCH LABORATORIES: LONDON, Eng., and WALKERVILLE, Ont. Manufacturing Chemists,

DETROIT, MICH.